

Small Tubes of  
Diatoms

Those in spirits or creosote  
water - for burning on covers or slides,  
and mounting in fluid are marked  
Crude.

5.

1 (Harwich W. K. Bridgen)  
*amphipora vitrea elegans* (var. scars), *Fusigella acuminata*, *Pleurosigma balticum* (large  
and small), *Stigosum*

2 (Harwich W. K. Bridgen)  
*Pleurosigma balticum* (large and small)  
*Stigosum*

3 (Hordon W. K. Bridgen)  
*Gamptylodiscus spiralis* (scars), *Cymatiphora*  
*torea*

3 Grimsby, Lincolnshire - J.N. (Feb 1882) 5, N. 328  
*Pleurosigma formosae*, *balticum* (large & small)  
*Stigosum*, *autumnale*, *undulatum*, *Nad. formosae*

5.

4/Harwich

W.H.Bridgeman

*Pleuronigma formosum*, *Hippocampus* (large  
and small). *Actinocyclus mediterraneus*

5

7/Harwich "Bl. 1" W.H.Bridgeman

*Pleuronigma hippocampus*, *angulatum*, *anophyllum*  
*oratum*, *Narcine pygmaea* (B. minutulus Wils.)  
*Stauroseris clinea*5/abendles - River Dee - D. dedform/  
*Synidium minutissimi*, granular? (After S. ci-  
-culans)

6/Harwich

W.H.Bridgeman

*Pleuronigma fasciatum*, *Hippocampus*, *angulatum*,  
*stigmosum* (scarus), *Surirella gemmea*8/Harwich "Bl. 2" W.H.Bridgeman  
*Tryphosella acuminata*, *gracilis* (empty)8<sup>2</sup>/Breydon, Norfolk R. M. Thw, Sept. 1 1881  
Camphyrn. <sup>calcareous?</sup> *helodes*, *Tryphosella*, *Cylindrus* (Hartmann)  
*Pleur. fasciatum*, *angulatum*, ~~straight~~ <sup>irregular</sup> ~~empty~~ <sup>empty</sup> ~~only for benthon~~  
— (area 123) *Baileya* (varia), *Spith. tuberosa*, *mentaria* var. *brevirostris*9/Flordom 1. W.H.Bridgeman  
Calyptidium costatum, *Narcine cuspidata*,  
sericea, *Pleuronigma tenerum*, attenuatum,  
*Lunaria diadema*, *N. tenuis*, *lineatum*, *tenuum*,  
*minutissimum*9<sup>2</sup>- Breydon Norfolk  
Calypt. clypeum — *new party* 1883

S.

7

- 10 (Ipswich - Wigham Jan N. K. Bridgem)  
*Pleurozium quadratum* (large and coarse  
 stratified)

- 11 (Formesby N. K. Bridgem)  
*Trichostomum exiguum*, *Adiantum trichoides*,  
*Navicula scutellariae*, *adnatae*: *Navicula* sp.,  
*Othismia punctata*, *Epipterygium gibbum*, *sorae*  
*Cambellia muculata*, *Cocconeum lanceolata*, *Lymnaea*  
*capitata*

- 12 (Harwick "H. no x. Freshwater" N. K. Bridgem)  
*Navicula cuspidata*, *amphibea*, *lemenii*?,  
*Nitzschia delicatula*, *Gymnophlebia solen*, *Pleuro-*  
*Zizanioides attenuatus*, *Pinnularia viridis*,  
*Serirella splendens*, *australis minutissima*?  
 ex. sp.

5

- 13 (Harwick "H. 3" N. K. Bridgem)  
*Pleurozium helictinum*, *stigodium*

- 14 (Harwick "H. 7" N. K. Bridgem)  
*Pleurozium formosum*, *stigodium*, *Bethia*  
 (large and small), *quadratum*. *Navicula lemenii*,  
(large and minute 1 mm)  
*conica*, *Lycophyllum decolor*, = *S. pulcherrima* (Bz)  
 a. *Phycomyces* *septata*, *alata*, *constricta*, *ampho-*  
*lycaline*!

- 15 (Hull. A. Harrison)  
*Pleurozium elongatum angustum*, *fasciolatum*,  
*Hypothecaria* sp., *Stigiloma* *nitidum*, *guttatum* B, *Liniella*  
lindbergii, *sp. regularis*, *maculata*, *aurata*, *adspersa*,  
*germanica*, *Taybl. punctata*, *Pinnularia* *perfracta*,  
*Schizogonium* *crus*, *Ruppia* *affinis*, *navicula*  
*peristicta*, *tumens*, *hastata*, *amphibia*,  
*dagffora*, *americana*

5

Hoxdon - "Bl. 6"

- 16 (Richmond, Virginia - April 1859 D. S. Gregor) *Cocconotus* *discrepans* *var. sp.* - *Ostrea* *bullocki*  
~~(soft-shelled marine)~~ *Pinnularia* *oblonga*, *P.*  
*tenella*:  +  *nudata*, *globoidea* W.G. *Navicula*  
*austriaca*, *gibberula* (= *limosa*) *Stauroneis* *gracilis*  
*analis* t-

- 17 (Felthorpe *M. Wigham* q:51)  
*Navicula* *sericans*! *Nitzschia* *curvula*! (medium  
 form. same as in 3.196, 5.197 - *Pinnularia* *gibbe-*

- 18 (Hull April 1859 D. H. Murray,  
*amphipod*, *helodus*, *Surirella*, *Brightwellia*, or other  
*striate*!, *amphion* *texana*, *salina*!, *Epithemia*  
~~extinct~~!, *Westermannia*!, <sup>and var.</sup> *Tyblionella* *gracilis*  
*Pinnularia* *radiosa*, *Navic.* *lunari*, *Cyclot.* *Kut-*  
*zingeri*, *Synedr.* *circularis* *gracilis*; *Velutina*  
*var. punctata*, *Velutina* *var. punctata*

See 5.174

5.

- 19 (Arthur's Seat, Hunterbog - *Grew. 33*)  
*Canthylocidium* *costatum*, *Nitzschia* *sigmoidea* Am.  
*Ahora* *ovata*, *minutissima*, *Denticula* *lineata*, *Epithemia*  
*superba*, *odontoides* = *parasiticum*, *Cornut* *flavus*,  
*elliptica*, *solea*

See 40, 5.30.

- 20 (Drain W. end of Duddington Lock 4:54. *Grew. 27*)  
<sup>digitata</sup> *Pinnularia* *oblonga*, *gracilis*? *acuta*, *major*, ad   
*Stauroneis* *gracilis*, *Dubia* W.G. - *Navicula* *sonori* (W.G.) *gib-*  
*lousa* (*limosa* Kutz) *dilecta* *hyalochroa* Mch. *ostend*. *richard-*  
*Cyclotilla* *speculifera*, *Gonfthemus* *integerrimus* W.G.


- 21 (Drain S. end of Duddington Lock 4:54 - *Grew. 29*)  
*Stauroneis* *analis*, *gracilis*, *achm* *anthedoniana*,  
*costatum* *modicum* *constructione*.

5.

- 22 (Hunterston, 2000 ft., near the gate in stone  
*odontidium mesodon*, *mutable*. Melviller  
 varian)

(for Crude see 439)

- 23 (Aberdeen, Pools near sea 19:45 h D. Roff)  
*Schizoneura dillwynii*:

frustules oblong elliptical  no thick  
 wall  with *Peronella* - Kutz  
 (for Crude see 425)

- 24 (River Dee, near Threave 22:45 h D. Roff)  
*Schizoneura dillwynii*: 

<sup>long</sup>  
 frustules oblong - at vise S. obtuse : no thick  
 other  with Nitze, until Melville  
 (for Crude see 424)

5.

- 25 (Isle of Wight)

*Cocconeis Grevillei*

25<sup>b</sup> same from a specimen in Provost Lab. 1790

- 25<sup>c</sup> (Roxburgh, Dr. Gledhill's Park, Henricy, Oct. 62) in cl.  
*Tetraclides emarginatus*, *Somphos. dichotoma* L. &  
*Cymbella v. sp.* - central cells

- 26 (Cardross, Neyland F. O. Keden OK. 233)  
*Berkeleya fragilis*

~~very long~~  
~~not thick~~  
~~posterior stalk - and out of specimen~~  
~~(for more detailed see 56 : )~~  
(Crude)

- 27 (New York, on *Giraultia american*)  
*Podocystis americana* Bail.

S.

28 (Torbay  
Lichenophor. flabellata)

(for crude from Wales - See 438)

29 (River Shannon 7:53. Desjardin Smith)  
Diatomea grande

29<sup>2</sup> Hull - build G.N. "7" Oct. 365<sup>h+l</sup>

Pinnularia gracilis Sm = Deltothrix Sm var = D. gracilis W.G.  
f. 30, 31 & 32. Melotricha Borneri ex

30 (Hunter's bog. art. Hens Leth - Grev.)  
amphore minutissima, odontidium parasiticum  
empty - for build See S. 19 & 40

30<sup>2</sup> Cumlaea in a canal R.H. 10:59  
Norvegia clarinella (Eng. var) Little Pennula  
and Nav. appendiculata (Eng. See S. 195 & S. 379<sup>2</sup>)  
Harrison. - same as Crude ~~for build~~ ~~for build~~  
in S. 110.

8-19)

S.

31 (Hull R. Harrison)  
odontidium Harrisii

crude (for build see 468 57)

32 (Granton Quarry 6:5:54 D. Griville)  
Pinnularia directa (very small), P. exante<sup>directa</sup> small  
Pleurosigma angulatum?

33 (Granton Quarry 6:5:54 D. Griville)  
Pleurosigma angulatum? (var.?) Pinnularia  
directa! ~~Spumiferus~~ ~~Spumiferus~~ ~~Spumiferus~~ ~~Spumiferus~~ ~~Spumiferus~~  
Synedra unisetigera? asterionella formosa? Am  
Ulothrix cincta

5

34 (Granton Quarry D. Guille 49)  
*Pinnularia Johnsonii* - directe

5.

37 (Granton Quarry D. Guille)  
*amphor. saline*, *amphipora pulchra* very  
 seen)

crude (for boiled see two last)

35 (Granton Quarry D. Guille)  
*amphor. saline* S.V.

38 (Granton Quarry D. Guille)  
*Melosira nummuloides* very fine

much boiled

36 (Granton Quarry D. Guille)  
*amphor. saline* mostly F.V.

(for crude see L13)

slightly boiled

39 (Watt's Lodge, Derbyshire - 4:5th D. Rufford)  
<sup>windblown water</sup>  
*Pinnularia gracilis*, *Nanula amphibora*,  
*producta*, *amphirrhynchus*, *opina*, *bivalvis*,  
*Cymbella amphiceraspis*, *mentula* - very dark

S.

40 (Girrock, by the sea side in a spring 17:6:54) ...  
*Nariae* *anophrynsis*, *brevula*, *gibberula*, *firma*,  
*anophyllum* *stellatum*, *Coccinea* *cyanifera* - *Stauroneura*,  
*gracilis*, *aniceps*, *linaria* *denticulata* *sinuata*. *Cyclotella*  
*Mastagloia*. *Smithii*

41 (Grantown Inveray 5:54 D. Graville) ...  
*Schizogonium* - *or homocladum* sp. (?) *both anophyllum*,  
*demissum* (*felix*), *anophyllum* *pulchrum*

slightly bristled (see 115#s 5.360)

42 (Loch Leden, near the outlet D. Gregory 5.1)  
*Cyclotella* *Rotula*, *athosae*, *punctata*?

S.

43 (Loch Leden, N. the outlet D. Gregory - 5.1)  
*Cyclotella* *rotula* (plentiful), *othosae* *punctata*;  
*Cocconeura* *cornuta* (shallow water), *Stauroneura*  
*stigmaria* W.S. - *Gomphosphaera* *stigmaria* W.S. - *Nariae*  
*stellaria* (very scarce), *Lacustris* W.S.

44 (Exmouth - misf Cutler A.)  
*Schizogonium* *bogatum* fid. W.S.  
 received under the name of "Nyctice?"  
 (misf for crude see 431)

45 (Exmouth - misf Cutler B)  
*Schizogonium* *molle* fid. W.S.

received as "S. Smithii"

S.

46 (Sidmouth Miss Butler,  
*Schizoneura Smithii*

received as *T. torquata*

47 (Granton Quarry 5.5h D. Griffith)  
*amphipor. contracta*

crude (for brtld, see next)

48 (Granton Quarry - 5.5h D. Griffith)  
*amphipor. contracta*

slightly boiled (for crudulat.)

S.

49 (Granton, in pool by sea side, D. Griffith) <sup>5.5h</sup>  
*Schizoneura Smithii*

50 (Granton Quarry D. Griffith)  
*Luridella fastigata*, nitreous lime!  
*Schizoneura conigerum*, *actinotus subcylind.*  
*Ovalular oblique*

51 (Hornby fountain, May, Lancash 5.5h C.T.)  
diatom elongated ♀ - *Encyonema pertusum*

crude (for brtld in 116 - for crude  
of *Encyonema* see 435)

S.

52 (Woolwich)

(J.T. Norman)

*Pleurozium Hippocratea*, ..... (part 33)Pinnal. Johnsonioides, Schizoma ovata, 67-  
closter <sup>(see margin)</sup> Kingianum, Dicranum fusiforme, D. gracile  
aculeatum 218mm

for Balsam

53 (Cumbrae)

*Cocconeis contellum*

only pt for Balsam

54 (Kingsgate, Tipton)

(D. Graville)

*Cocconeis quadriformis* Juss.

for crude in 104)

52<sup>2</sup>. River Ape - Little part of S. 25<sup>2</sup>  
at mouth of river. Grafton station &

S.

55 (Woolwich 17:6:54)

(J.T. Norman)

*Pleurozium Hippocratea*, ~~Smilium laeve~~  
Nitzechia fasciata

(part. P. Littoralis)

(for crude see next)

56 (Woolwich

17:6:54 J.T. Norman)

Nitzechia fasciata - *Pleurozium Hippocratea*.

crude (for crude in 5.55)

57 (Albionstone

(imp. P. Hodges)

*Fragilaria capucina*, Melobesia Nation  
odontidea mesodon

Crude

5

5.8 (Fell End near Lancaster 6:6:5h C. 5)  
*Adontidium mesostolon*! *Fragilaria*. *versicolor*  
*(Scum)*, *Suriella litoralis* (Scum) *Admontidium fusiforme*  
*al. t. major* (Smith Small). *Syndra ulnae*: *collotonum*  
*vulgaris* (like *S. crispin*, but coarsened borders)  
 (for crude see 409)

5.9 (Pentland Hills 5:5h - Dr. Griville 59)  
~~*Epihemic gibba*, *sphaerica*!~~ *Denticula linearis*!  
~~*Cocconeis Thwaitesian*, *Navig. angustata*!~~ (Smith - H.  
*cervicula*! *sid.*) *Mastogloia Griville*! *Pinnularia*  
*hemitropa*. *Actinopeltis* *Aspera*!

6.0 (Tantallon, Berwickshire - Dr. W. Gregory)  
~~*Suriella crumosa*~~ *detrib.* - *Sp.*

7. 59<sup>2</sup> Warrilee river, Australia J.J. Norm 1861 no 3.a  
*Syndra ulnae*, *radicans* - for another sample from  
 same place see 5.485\*

5

6.1 (Granton Quarry 3:6:5h Griville 61)  
*Schizotrichia Grivillei*

See 88.

6.2 (Pentland Hills 5:5h Griville n. 58)  
*Gymbella maculata*! and *bocconea cyathiformis*  
 ("Gymbella very unlike any of Smith's" A.K.G.)  
 \* in *C. maculata*, Smith's show, but not of his figures

6.3 (Cumbrae 9:53 O. Kennedy)  
*Melosira Bonari*

(For crude, see 412)

5.

- 64 (Spearham Lurry D'Urville Jan 1854.  
*Schizoneura evanescens* Desv. (a moulana) = *S. dilatata*?  
*Podothecium* Schr. - (Throne one day)  
 64<sup>2</sup> High Bridge Yorksh. OK. 5.20<sup>2</sup>  
*Diatoma vulgaris* Thott and Houtt

- 65 (Exmouth - mis Butter C. )  
*Schizoneura Smithii*

- 66 (Peruvian Guano - Dr. Rooney)  
 bad - removed

- 66<sup>2</sup> Minatitlán near Vera Cruz G. Norm. & N. 343  
*Lepidolema murica*  
 fruct. of sand

5.

- 67 (Hamilton 12: 6: 54 Army N.Y. file  
~~Cyathula armata~~ (Lam.) Rommelsb.  
*Githemis turgida*, gibb., *Pinnularia major*

- 67<sup>2</sup> Edgefield Holt, Norfolk. - M. Wigham, Brit. Kitten  
 1862  
*Narcissus tenuissimus* (L.) 17. - Hibbert. undulata

- 68 (Stone in Stream from Carlton Park 6:54)  
*Zonophorina olivaceum*, *Lerinella minutula*,

- 69 (Lafayette, on the boards of a shire 16:6:54)  
*aetnæ austriacum lineare*. <sup>Spec. no. 77</sup>

5

- 70 (Hampstead - June 1854). Norm. no. 2  
*Navicula firma* (Smell) Lindbergh  
 (Smell)

for large see 5, 76

- 71 (Woolwich 17: June 1854). Norm. no. 3  
*Diatom elongatum* vulgaris, St. Louis  
 linear

- 71<sup>2</sup> Ellerker Lock, Hull. - S. Norm. 353  
*Gomphonema curvatum* m. 3, 73<sup>2</sup>

- 72 (Pentland Hills 5: 14 D. Gerrill 1857)  
*Odontidium heemati* var. for W. Sm.  
 (od. glaciella Nutz?)

5

- 73 (Deposit from ~~Hengay~~<sup>part of Hengay</sup>, above the Cliff - C. major  
*Pinnularia viridula* chatty, C. major  
 73<sup>2</sup> Brough near Hull - S. Norm. 1861 - S. N. 349  
*Gomphonema curvatum*

- 74 (Hull, ~~sold~~ <sup>and the light parts of</sup> 23)  
~~Tessellaria apiculata~~ <sup>apiculata</sup> M. J. S. - Nutz. Norm. Pleura.  
 angularis, fasciata, elongata, - all scarce)

- 75 (Woolwich June 1854 I. Norm. no 71  
~~Nitella aciculata~~, Pleurosigma slightly  
 fasciata elongata, Hippocratea (small & leaf)  
 Navicula ~~exigua~~ us, cryptothelus, Tryblia  
 -ella marginata, Cyclotella Kutzner p. 3  
 75<sup>2</sup> Scotter Bog near Hull - S. Norm. 396  
 Stauros Baileyana - Nutz. Lymnaea P. Sm  
 (three each)

5

76/Hampstead. April 1864 J.T. Norman No 17  
*Spirirella bisinuata* & *splendida*, N. sp. 2 sp.  
*modica*, *Navicula firmula*, *Orimula viridis*,  
*nobilis* - all large & firm  
 n.s.

See S. 70

77/Lafswede, in gravelly pool, D. Saville )  
 Nitro. Shi. aciculare, N. curvula, Navicula affinis, *Pinnularia viridis* mesostigma,  
*Stauronea granularis*, *Cymbellula affinis*, <sup>and Maculata</sup>  
*Spirirella linearia* (Scam.)

78/Bohemia, near Egr. J.T. Norm.)  
*Conopylepsidium clypeum*, *Navicula tumens*,  
*N. culta*, *Pyrenula bohemica* of Thunberg (Nikolj)  
*Penicillium hyalynella* Chr. (not Seg.) - in the water X. I.

not same as S. 215

N.B. my fish 78 got injured by a miliea godonii Marcelli  
 from Hull - Then he - a later birth and marked 70

5.

79/Teignmouth <sup>Brackish</sup>  
*Grammatophor. marine*? *Phallidium*  
*areolatum*-  
 The form of *G. marine* has since easily been seen by naked  
 eye in back  
 N. 3 - and much coarser than in S. 539 - Lemma ~~7. 539~~

80/Tilbury fort

amphiprot. alata-

80<sup>2</sup>/Sutton-hedge, Lincolnshire. G. Norm. 1862 No. 396  
*Spirirella nobilis*, *conspicua*, *costata*, *Stauronea*, *penicillata*,  
 Martogli. S. m. the var. - Nav. bailey <sup>of</sup> *Spira*  
 var. *lutea*, *Epith. argentea* & *Obtusata* ~~obtusata~~ <sup>obtusata</sup>

81/Tilbury fort brid.

amphiprot. alata - *Pinnularia cyprina*,*Tyblionella acuminata*, *Navicula convexa*,N. t. L. *Ligula amphiprot. rotunda* (new comb)  
<sup>Galeolaria</sup> <sup>Small (S. 527)</sup> (S. 540)  
<sup>(Cyclotella</sup> <sup>Thuringiana</sup> <sup>angustata</sup> <sup>and</sup> <sup>prolata</sup>)

~~Stauronea~~ <sup>the</sup> *Penicillium*, *Dactyon*, *amphiprot. new*  
 (Scam.)

5.

82 (Tillney foot                    J. T. Norman,  
amphoraeline ad: a. tenera

83 (Hull -                  Linn 1854 R. Harris) \*  
Tryblionella punctat., marginata,  
achmantus brevipes. Synder. affinis. Podo  
site montagnes &c

84 (S. Helens, Lersey, W. Dohr Jr. Landsp.) \*  
Isthmia enervis

84<sup>2</sup>. Dolgelly Earth - Light earth - for heavy seas 120  
Eunotia diadema

Bread in turpentine/sea  
in spirit 30)

5.

85 (S. Helens, Lersey, W. Dohr Jr. Landsp.)  
Isthmia enervis

Bread in acid

86 (Powburn <sup>in a bog</sup> mouth, near Prestwich, April 8:7:54)  
Pinnularia. i integra. viridis

87 (Yarmouth                    R. Figham N. 1)  
Maricula elegans, Pitmeule peregrina, th

S.

88 (Yarmouth A. Wigham No 4)  
*Cantharidium ciliatum* (very scarce) *Seri-*  
*nella striatula*, *gemma*, *fastuosa*, *Pleurodig-*  
*angulatum*, *elongatum* (<sup>Hypothecaria?</sup> *acuminatum*), *Pinnular-*  
*cyprius*, *Navicula elegans* & *peniculus* (both rare)  
~~very very few~~  
*Tuberculina marginata*, *punctata*

89 (Breydon, Norfolk - M. R. Wigham No 3)  
*Serulae arcuatae* (<sup>(= striatula small)</sup> *Smilium*), *gemma*, *constricta*, *Tegulina*  
*marginalis*, *scutella* (very scarce) *Actin. undulata*, *Eupodium*  
*fulvum* (*scrofa*) *Canthar. ciliatum*, *Podos. maculata* *Brid.*  
~~repudians~~ *repudians* *F. V.*  
*dulcis*, *ochracea* & *Smilium* (very scarce). *Thomatis*, *Navicula*  
*smithii* *Brid.* - *Cyclot. peniculus*? *inflata* *leptostoma*

90 (Moanarone, Co. Cork - April 5. Prof. Smith)  
*Meridion constrictum*, *Fragilaria ericetorum*,  
*orthotilis orichalcea*, *odontidea mesodora*  
*Navicula chorobridea* & (*N. pusilla* or *torquata*?  
 perhaps mixed from S 247)

The orthotilis should be mounted dry

S.

91 (Falls of Clyde 1:7:5h)  
*Epihtemica ocellata*, *longicornis*, *gibba*.  
*Cocconeis Thwaitesii*, *Denticula sinuata*

92 (Arran. 2 m. N. of Brodick 11:7:5h)  
~~Navicularia pinnaria~~, *rhomboides*, *seriosa*:  
*Pinnularia hemiptera*, *ruvidii*, *stauronea*  
*gracilis*, *Nitzschia curvula*

92.1 Anchor. in St. George Bay, Bermuda 40 to  
 centrl. notates, the two (modestus green) *Strophidium* <sup>11:7:62</sup>  
 flatte or rotta frises, crustose. Eth. N.Y. London form. *Vener*  
*tanakae*, *Eupodium* *Brid.*, *Pleurodictyon* *Brid.* *Velutina* *Joh.*

93 (Brackish pool at Corrygheilly 10:7:5h)  
*Navicula elegans*, *amphibaea*? *gemma-*  
*rosa*, *punctulata*, *stauronea crucicula*?

Dried up & removed  
 apparently very bad

5

94 (Goatfell, Aran. off 27° 50' N. Lat. 11° 54' E.)  
*Navicula rhomboides*, *crispinaria*, *serians*  
*Pinnularia biceps* (interv. var) *vividis*, *rigida*,  
*Eunotia bidentata*, *tricodon*, *Cymbella scutella*  
 (Scen.) *Himant. major* (?) *ment.*, *l. rot.*)

95 (Paw-burn mouth, near Portwick; same bog as 86).  
*Pinnularia vividis*, *nobilis*, *affinis*. *Surirella*  
*brevata*, *Mitella dubia*, *Stauroneus gracilis*

96 (Banks of the Sunnill among Sphagnum A.K.G. no. 83)  
*Pinnularia hemiptera*, *vividis*, *borealis*, *gra-*  
*-ciliata*.

96<sup>2</sup>  
 [mixed in a little of Melville's nummuloides from  
 Constant Quarry - which outcrop looks down upon the  
 water to Sunnilock little]

5.

97 (Woolwich J.T. Norman)  
*Surirella striatula*, *Pleurostigma angulatum*,  
 (Smell), *fasciola*, *Amphipora alata*, *Stauroneus*  
*clina*, *Typhlonella angustata*, *Mitella dubia*,  
*Ctenium*, *muscorum* K?

98 (Super coast - H. Smith, & Oct. 27 )  
*Mastogilia lanceolata*, *amphora affinis*, *Stau-*  
*-roneus salina*, *Pinnularia*, *Cyprinum*, *Mitella*  
*obtusa*?

99 (Cromer, Fresh water A. Wigham no. 5)  
*Campylodiscus costatus* (Scen.), *Opercula* (very  
 Scen.) *bymatophorum*, *Solea* (common), *Navicula*  
*amphibia*, *euphiolata*, *Amphora ovalis*.

96<sup>2</sup>? Rame, on <sup>the</sup> northern stream, F. Other 12:3.63 - old Sh.  
 "odontidea" Nuttall? "D.N." = denticle inflata

S.

100 (Wells, Norfolk, brackishwater R. Wigham No. 6).  
*Pleurosigma fasciola*. *Tyblionella gracilis* New  
 reginae? *Pleurosigma* ~~turneri~~ <sup>Turneri?</sup> Nitzei  
 dulcis. *Navicula cuspida*, *tumens*, *americana*,  
*bicornuta*, *solen*. *Collectaria elegans* <sup>exigua</sup> *fuscula*.

101 (Holkham Gap, Brackishwater R. Wigham No. 7).  
*Podosira montagui*, *maculata*. *Navicula didyma*,  
*liber*, *elliptica*, *lunaris*, *granulata* (Koenig), *Tybl*  
*onella marginata*, *punctata*, *accuminata*. *Pleurosigma*  
*angulatum*, *stygium*, *Hippoc.*, *ostuarius*, *Litt. gemma*, *douglasii*  
*amplic*, *Staur. salina*, *Nitz.* and *M. othos. mariz.* *Euped. scutellari*!  
~~actinocyclus undulatus~~, *Canthyl. punctata*,  
*Cocciundicium minus* Nutz. *Mastogl.*

102 (Felixstow, Norfolk R. Wigham No. 8)  
 " *Surirella striatula*, *Brightwellii*, *festuosa*,  
*Navicula orafinum*, *punctata*, *tumens*, *brevirostris*  
*Tyblionella acuminata*

S.

103 (Harwich, Brackishwater R. Wigham No. 9)  
*Pleurosigma elongatum*, *fasciola*? *Hippocampus*!  
<sup>annulus</sup> <sup>subcuneiformis</sup> <sup>obtuse</sup> *Tyblionella apiculata*? *Nitz. dulcis*, *psammophila*? *S. marginata*,  
*Surirella striatula*, *Stauroneis salina*, \*  
*Mastogloia Smithii*. *Cyclistella kutzingeriana*  
 & this must be distinct - see S. <sup>380</sup> 5-697, 296, and 520-2.

104 (Hornung, Norfolk R. Wigham No. 10).  
*byssella Threnbergii*!, *Odontidium Tabellare*!  
*Cocconeis lanceolata*, *Surirella bideriata*?  
*amphora ovalis*, *Pleurosigma attenuatum*, *Cymatopera elliptica*.

105 (Donaghadee R. Kennedy July 1854)  
*Mastogloia apiculata* No., *Navicula*  
*helpebralis*, *Stauroneis pulchella*, *Podosira maculata*,  
*Canthylodiscus Ralfsi*, *Surirella lata*, *Pupillaria pulchra*,  
*Cocciundicium radiatum*, *Actinia* (this large per. actino-  
 -ycles undulatus, *Grammatophae Serpentina*, *Pleuro-*  
*sigma fasciola*, *elongatum*, *Schizoporella cruciger*.

5

106 (Drain, below reservoir, Donally, Perthshire, R.K.G. No. 85)  
*Dontidium mesodon* var (*od. glaciata* Kutz.)  
 same as in 5.72 - Canaliculi few and not  
 continued (the end)

107 (Bog at Bonkield house, Perthshire 7:54 R.K.G. No. 88.)  
*Navicula* & homobidae, *craibi* novia, *Leptians*, *Dontidium*  
*mutabile*, *Harringtoni* (rare), *Pinnularia stauroneiformis*,  
 interrupt. (stem) *vividus*, *major*, *Hemimelania* *dilatata*,  
*arcuata*, *pectinata*, *Nitzschia curvula*, *Eunotia tetraodon*,  
~~metabeta~~ *nivalis*! *Spitheria rugosa*

108 (Side of the Tummell among Sphagnum, 7:54 R.K.G. No. 89)  
*Pinnularia borealis*, *lata*, *vividus*, *major*,  
*Lemniola densitia* *tetraodon*.

108<sup>2</sup>. Thame, Oxfordshire. 7.54 R.K.G. No. 89

*Cymbella affinis*

5

109 (Spring in a Fir wood about 1500 ft. E. of Hilli-  
 Krautlie 7:54 R.K.G. No. 91)  
*Poecconeis Thwaitesii*, *Navicula firma*, *cocconeiformis*,  
*Surirella heteriata*, *amphioxys*, *linearis*, *Pinnularia*  
*vividus*, divergens nov? *Stauroneis granulata*, *Pleonicula*  
*sparsa*, *Cymbella affinis*, *obtusa*, *Gonophorina clavata*

110 (Wells, Norfolk. R. Higham No. 2)  
*Microcystis trigonum*, *navicula liber*,  
 (scarce) *Pinnularia cyprinea*, Nat<sup>l</sup>: *clathrata*  
 Eng<sup>l</sup>: *M. cruciformis* Drak<sup>l</sup>?

111 (Corrieghills, Aran, between the cales and  
 salt marsh 10:5:54)  
*Cymbella cuspidata*, *amphora ovalis*, *Cyma-*  
*toflella elliptica*, *Solen*. *Dontidium mutabile*

S.

112 (Cumbrae in deep S.W. pools near the sea

S. of Millport 9:8:54

*Epihemic longicornu*, <sup>argus</sup> ~~acutum~~, Mastaglina  
Smithii : (partibus suberis)

113 (Cumbrae in a ditch S.W. of village 9:8:54)

*Campylodiscus costatus*, *Ptilialis (very rare) atheros*,  
*arenaria*, *Amphora ovalis* (amphor-minutissima)  
and a doubtful parasite observed before being cleaned) *Cymo-*  
*Tfleur-solea*, *elliptica*

114 (Corriegeoffs, area in the salt marsh 10:8:54 -

on *Bladophorus flexuosus*)*Mastaglina lanceolata*, *Achnanthus brevifolia*,  
*Subsepsidium*, *Epihemic muscularis*, *Cyclotella*  
~~sp. obs. minor~~  
~~sp. (?) tenuis~~ ~~thin~~ ~~that~~ ~~widestard~~

S.

115 (Ulverstone, Salthouse Aug. 1854 Mif P. Noddy no 60)

*Pinnularia Tabellaric*, <sup>major, notula</sup> *gibba*? *Tegularia*,  
*Wiescens*! (<sup>some</sup> *in threads*)

116 (Heswick Tarn, Ulverstone, Aug 1854. Mif E.H. no 35)

*Gymbella strenuosa*, *affinis*, *Nanula spheen-*  
*thon*!, *cuspitata*, *temer*? (Same as in Lang-  
muir), *infusa* (very scarce), *Pinnularia nobilis* (no

117 (Salthouse, Ulverstone - Aug. 1854. Mif P.H. no 60)

same as S. 115, but now boiled and the  
*Pinnularia* better for balsam

for crude see 416

5.

- 118 (Brodick bay in the burn - Aug 1852 )  
*Dasyphora* *Brockii* ! Lynden Pritchell.  
 (Same as 17, but in the more of the *Dasyphora*)

119 (Mullock Craighead, Carrick, D. Landsborough 22.8.53)  
*orthosira* *orichalcea*

120 (Dolghelly deposit - C.T. calcined)  
*Seriella notabilis* - for light see 5.84<sup>2</sup>  
*Pinn. acrosphearia* Sm (as to twice only = *P. major*  
 var?)

5.

- 121 (Lough Mourne Deposit, A. Kennedy 7.53.)  
 Heavier diatoms  
*Cyclotella minutea* Nutz / *C. antiqua* Sm. n. sp.,  
*Staurosira acuta*, *Phaeocystis*, *gracilis* K.

122 (Lough Mourne Deposit. A. Kennedy 7.53)  
 Lighter diatoms  
*Cyclotella minutea* Nutz (as in 5.121). ravig.  
 .... like N. obesa but much smaller and very diff.

123 (Umbrav, Fresh water, W. the sea, N. of Millport  
 - Aug 1854. Lighter parts of 139)  
*Mastogloia* Smith, *Staurosira punctata*, [Phi.  
*Nemicia ocellata*, ~~granulata~~, gibba - all species]

for crude see 1.14

5.

- 124 Ditch between the two water reservoirs above Adens  
Aug. - 22:8:54) *Pinnularia viridis* fm

- 125 (Sufex 25:8:54 - W. Smith)  
*Pleurosigme macrum*! and *Fasciola*  
(*l. M. Scarr*), *Tryblionella scutellum*

- 126 (Lewes, Sufex 25.8:54 W. Smith)  
*Epihemicia dores*

5.

- 127 (Lewes, Sufex. 25:8:54 W. W. Smith)  
*Epihemicia turgescens*

- 128 (Premnay peat after heavy falls, mixed with  
the very heaviest of S. 129. See Annals of Nat.  
Hist. 1848 p. 94 - also here are) *Narciella lemen*.  
*Pinnularia alpina* (var) borealis (var) lat (very rare)  
- - B. P. L. monodonta

(accidentally mixed with this a rock tree w. <sup>found</sup> different  
kind of bark)

- 129 (Premnay peat after less burnt with the cleanest  
of S. 128)

*Pinnularia acuminata* (but often blunt)  
~~and~~ *Pinnularia alpina* (few)  
*Oxav. lemen* (very scarce). *Pinn. hem. lter.*; *des-*  
*tylus*, *Eurosthem. Hoch*; - *Polygonum* (?) (very scarce)  
*Staur. acuta* - *Luz. bilobata*, *nobilis*

This contains nearly all that is good in S. 130  
+ the *Pinnul.* chiefly differ from *P. viridis* by three ss  
in O. O.

130 (Prennay, perhaps cleaned - Lightfoot 95.128, 129)  
orthosia orichalcea chiefly

131 (Rock, mouth of Tamar 8:54. G. & Densley)  
"Cocconemis cistula and several crista-  
tella, neither typical forms" Dr. Sm.  
Dentical. ~~obtusa~~ Dr. obtusa var. ~~obtusa~~


132 (Bræmar 8:54 "Dr. Balfour") - R.K.G. 106  
Himantodera major (fid. W. Sm.) Pectinal,  
Punctaria 3-odor, Pinnularia viridis & a. sp.  
alpina (Scen) Nav. firma, confertioria,  
Mallotis nivalis, odontidium mesodon

5.

133 (Bræmar "8:54 Dr. Balfour" R.K.G. 107)  
Tabularia fenestrata, and flocculosa (var. ventricosa -  
rare - Scen) Odontidium mesodon, Synechia  
longissima? (varicosa var. pedicellata?)

134 (Glen-Tilt 8:54 "Dr. Balfour" R.K.G. 111)  
Tabularia flocculosa var. ventricosa (abundant),  
fenestrata (Scen)  
alluvia. "

135 (Ballochian 8:54 "Dr. Balfour" R.K.G. 112)  
Cymbella setosa (abundant). Cocconeum cyprin-  
iforme, <sup>infusorij</sup> (Tabularia) <sup>infusorij</sup> (Tabularia)  
Dentical. Setaea, tenuis, Cocconeum  
fuscatum, Gonfionem diastoma (abu-  
ndant) Nav. tenuis & th. rugosa - Synechia



- see 3.392

5.

136 (Pm. Macdui - 8:54. D. Balfour. A.K.G. n. 113).  
*Pinnularia*, *lata*, *long*, *rav*, *lunata*, *monodonta*,  
*tetragon*, *triadon*, *Melosira*, *midulis* (◎) *Nimantid*.  
*revera*, *Tabellaria* *flocculosa* *var* *ventricosa*, *Cyano*-  
*bella* *exigua*, *Achnanthidium* *lanceolatum*

137 (Caullochan 8:54. D. Balfour. A.K.G. n. 114)  
*Diatomella Balfouriana*, *Synodi*.

138 (Breamar 8:54 D. Balfour. A.K.G. n. 115).  
*Lunula* *bidentata*, *triadon*, *narcissus*  
*semis* ?, *rhomboides*, *crescens*.

5.

139 (Breamar 8:54 D. Balfour. A.K.G. n. 116)  
*Melosira dibans* Kutz ! *adrentidium anomatum*  
*Im.*, *odonthyridium* (*sporangia*) *parvum* (11111)  
*and med. don*

140 (Breamar 8:54 D. Balfour. A.K.G. n. 117)  
*Pinnularia*, *stauroneiformis* (*long*) *alpina*, *lato*,  
*acuminata* (*rav*) *astertonella*, *Ropeiformis*,  
*Gonophorema capitatum* (?) *fidei W.S.* *narcissus*-  
*rhomboides*, *crescens*, *coccoidea* ! *Lunula*,  
*lunata*, *lineata*, *Lunula* *mediana*, *tetragon*,

141 (Breamar 8:54 - D. Balfour A.K.G. n. 120)  
*Narcissus* *coccoidea*, *crescens* ? *Tabellaria*  
*flocculosa* *var* (?) *Nimantidium bidentata*, *Pin-*  
*-nularia*, *borealis*, *Lunula*, *triadon*, *gracilis*,  
*nicilis* W.G., *Melosira* - *nivalis*, *Cymbella*  
*lunata* W.S.

\* true, with a conut - fidei W.S.

5.

142 (Bræmer 8:5h D. Balfour R.K.G. n.129)  
*Melosira* <sup>*nivalis* (Horn)</sup> ~~*distans*~~ orthos. spinos. odontidea  
 mesodora, *Pinnularia* lat., *Himantidea* bidens  
 Nav. angust. tr. (metres recently a Cymbell) Cymbell. setosa  
 var obtus. affinis

143 (Bræmer 8:5h - D. Balfour - R.K.G. 125)  
*Tabularia* fenestrata, flocculosa. Nav. clombria,  
 ex spinosa, *Melosira* distans  
*Himantidea* bidens, *Syndesmion*, *odontidea*  
 mesodora, *Eunotia* tenuis, *guttulata*, *bidentata*  
 , nav. leucostoma, *Cymbella* ventricosa?

144 (Gerbaldado - 8:5h D. Balfour R.K.G. n.127)  
*Denticula* obtusa

5.

145 (Sunbrick near limestone, 9:5h Mif. P.A. n.111)  
*Pinnularia* radiata, *Nav.*

146 (Kent map. Alvorstone, 9:54 - Mif. P. Augr. No 4)  
*Pinnularia* acutipennis, *Lemniplana*? Nitrodoa line-  
 aria, *Surirella amphioxys* var. *Cyclotella* minutula  
 Kutz (very small) *Stauroneis punctata* (very small) gracilis,  
*Staur. acuta*, *cruciata*, *navic.* *leveiformis*,  
*Cymbella* affinis, *Himantidea* (very small)

147 (Hackney, Liver 1851 I. T. Norman)  
*Fragilaria* capucina, *Cocconeis* pediculus,  
*adnathera* etili

> 146<sup>2</sup> / Pond near old museum in Kew Garden. G. Norm  
 1861 - 8.V.326)

*Denticula* decipiens (parasitic - Nav. cryptopeltis  
 formula 405<sup>2</sup> - all 5.153

S.

148 | Salthouse Sands (Ulverstone 9:5h. Miss H. 16)  
Nitzschia digma K.

149 (Trelawny in Scam 10:8:5h N. Wigham.)  
Balticum and Balticum  
Pleurosigma littoralis, Trigodon, quadratum (Scam)  
accumulatum?  
+ Coscinodiscus minus Kutz. Dujon  
australis, Navicula punctata, curva, tenuissima,  
ampora affinis utra, Orthotrich. marina. (Trippel  
ampora affinis, Nit. spudata  
inexplicata, Tresch. altern. Exop. conf., radiata,  
+ Complanula parallela - alvearia Scam)  
Leptophidium parvulum - aleuticum Scam

150 (Same as last, but the light parts)  
Pleur. littoralis, Trigodon, accumulata,  
Balticum, quadratum, &c.

S.

151 (Bramcaster near Titchwell, Norfolk. J. Brightwell)  
Surirella Brightwellii, evat var, Pleuro-  
sigma quadratum, Melosira nummuloides,  
Stauroneis crucicula. Nov. brevi w.gr

152 (Spurwick R. Wigham - Heavy parts)  
Campylopedion Hodgsoni, Pleurosigma stigia,  
decorum  
? Accumulatum fuscum, Balticum, <sup>Wm</sup> Bosenia-  
tissimum, montagnae, <sup>Wm</sup> granularis  
decum or culicaceum - Didduckhe. Lombus  
forte  
(very scaly) " Treacleallum ~~Hodgsoni~~ <sup>Wm</sup>  
Habil. L. 75, conwaya, Wester. in Nit. angulosa  
+ actinopelt. 16-mm. ricer. striolata (Hordeum) Epith.  
maculata (dotted form) <sup>Wm</sup> Marinum form  
153 Same as last - but in lighter  
light parts  
very bad

153 (udder of whale, "Towlow of Hull" in Melville Bay)  
N. Lat.  $76^{\circ} 2'$  - Bathy of  $42^{\circ}$  and  $42^{\circ} 2'$  - G.N. 363  
Leptophidium gerillae, Sigmoidea - Sphaerula

5.

- 154 (Redesdale - A.K.G 144)  
*Narcissus pseudonarcissus?*  
 The greater part is narrow - and is perhaps Coll.  
*vulgaris*.

This is said by Dr. Griseille to have been  
 in tuber and then a Colletome

- 155 (Breydon, Norfolk W. Wigham)  
*Spirælla strigilata* chiefly

- 155<sup>2</sup> Yearmouth R. Wigham no 7 Kellie 1862  
 (var. ~~multiflora~~, Not ligulata / trichia) In tub. quadrata - Pleur. bifida,  
 lam. ~~gathered~~ - as nest. but the light  
 parts separated from the

- 156 (Breydon, Norfolk R. Wigham)  
*Pleurozium elongatum*. Strigilis.  
*Spirælla strigilata*, Tayloriell marginata,

- 156<sup>2</sup> Breydon. R. Wigham no 7 "for F. Kellie 1862  
*Netiopodelia* sp? (Kellie's fig. bl. decumbens) ligulata  
 Nut grass. Pleur. strigilis, elongata & Sur. strigilis

5.

- 157 (Broadwater, near Chalk, Kent. M.J. Norman)  
*Pleurozium intermedium* - (also *P. quadrata*,  
*bifidum*, *fasciolatum* & *scorpioides*)

- 158 (Cupfield, I. 10:54 J.T. Norman)  
*Eugeonema prostratum*

in upper Govd  
crude. (In tubers)

- 159 (Pond between Kandersteg and Schwarzenbach,  
 Switzerland. July 1854 Old C. O. Miles)  
*Cyathella truncata*, ventricosa, coccinea  
*equiforme*, *spithamea*, ocellata gibba  
 narcissus

Died up & remastered (not very good)

5.

160 (Kandersteg, Swiss Revd C.P. miln 1854)  
*Epiptemic ocellata*, *Corbetti*, *Sectice*?  
*adontidens*, *mesodens*? (*Dentice inflata*?)  
*Gomphonema dichotomum*

161 (Birdshope crag, Redesdale A.K.G. 136)  
*Nauicula periflamea* Kutz.

162 (Redesdale A.K.G. n. 138)  
*amphiflora*, *pellucida*

5.

163 (Redesdale A.K.G. n. 139)

*Lunaria gracilis* Or.

164 (Redesdale A.K.G. n. 141)

(~~adontale obtusa~~), *Gomphonema texanum*  
*Lymnaea oxyrhyncha*? (*holmae* & *tenuis* Kutz et S. Longf  
 var) - *diform elongatum* var. *dentatum*

165 (30 miles below London - I.T. Norman)  
*Pleurosigma fasciata* (Stone & Lept)  
*angulatum*, *amorphoides*.

S.

- 166 (Ponds near Haverfordwest, T. O'Brien on 63  
*Synedra radicans* (solitary products plentiful)  
 [lanceolate ovate, and ovate large (S. crumosa var?)  
angusta, *Cyclotella punctulata* var]  
 Medium density of S. 188 and S. 189

- 167 (Hull R. Harrison)  
*Pleurosigma subcrenatum*, angular, entire  
 - median - Schizogonium crucigerum - and Myrion  
 contracta, Baileya paradoxa var

- 168 (Inverkip glen, Sept 1854 A. Kennedy)  
*Cocconeis Thwaitesii*, *Denticula tenuis*,  
*Cyathella scotica*?

S.

- 169 (Deposit from Lord Roden property Co. Down)  
*Navicula sericans* (var. punctata Shantz)  
*Himantodora undulatum* (Benthic nodose Shantz)

Lame on Mourne mountain 2 yds away

- 170 (Lillhaggssjön, near Åre, Sweden, J. T. N. Dahl)  
*Pinnularia lutea*, *Coscinia radiata* (Mittess-nick),  
*Navicula leucostoma*, *Benthic. monodon*, *nicodina*, *diodes*,  
*3-don*, *leptodon*, *guineensis*, *diodes*, *8-don*, *goda*,  
*10-don*, *terra* (all scarce but those underlined)

- 171 (Christiania, Norway - deposit J. T. Nissen)  
*Melosira nivalis*, *Fragilaria cordata*

5

172 (Santa Fiore deposit. Heavier parts C.T.)

5.

173 (Deposit from Holderness R. Harrington)

173 (Santa Fiore deposit. Lighter parts C.T.)174 (Hull. R. Harrington - Dec 1854)  
~~Epi~~<sup>Stictic</sup>~~hemic~~<sup>Westermark von</sup> ~~americana~~ Mastogloia Smithii,  
Synedra tabularis slender and long (with 1  
long stem) - affinis (blended and trouble with S. tenuis)

Dec 1854

176 (Redesdale 8:54 A.K.G. no 142)  
Denticula tenuis, sinuate - Synthecium  
sticticum (in thick fol.) - <sup>Coral. helveticum?</sup>  
(or wt. corall. epiphyta var) C. scoticum, C. cornic. circinata, <sup>lunula</sup>  
two have been with my 8, and is therefor D. infl. var177 (Redesdale 8:54 A.K.G. no 143)  
Synthecium constrictum; ad G. acumin-  
natum (Scarce). <sup>var.</sup>

X

(Drudup & <sup>reputedly</sup> water)  
Parsim and dry ??  
Is this their way ??

5.

178 (Clay, Swansea Dock M. Okeden )  
*Epithemia muscularis* with moniliform trace  
 and another with faint trace (possibly described by Smeath)  
 and which is probably *E. weddelliana* of Hertz  
 pressed up & algae mounted

179 (clay, Brickyard, Carmarthen M. Okeden )  
*Saccostium Favus*

180 (clay, Brickyard, Carmarthen M. Okeden )  
*Didymella clombus*, sp. (see 5. 187)  
*actinocycles sedenarius* var.

5.

181 (Haverfordwest, T. Okeden - OK. 17)  
*Lurinella apiculata* N. Sm (or var. *s. minuta*)  
 mud of & algae mounted

182 (Haverfordwest T. Okeden OK. 75)  
*Cocconeema cistularia* Sabellari: penetrante

183 (Milldam, Haverfordwest, T. Okeden OK. 71) partly  
*Gomphonema generatum*? *Sphaerula*  
*lacustris* (very scarce)

S.

- 184 (church land, 7 m N. of Haverfordwest OK. 123 & 124)  
*Navicula firma* var! *Stauroneis gracilis*?  
*Hemantidium undulatum*, *Pinnularia acrostica* - (Pim. bicolor w. g. said thicker, but scarce?)

For heavy parts see 310.

- 185 (Spring near Haverfordwest. OK. 39)  
*Navicula crassirioric*  
 Scum pink & damp in sun (say)  
 on Collier, vulgar?

- 186 (Swansea Dock - OK. 76)  
*Amphora salina* (too much boiled)  
*Surirella ovata*, *Rhytidionell* -  
 (Nitz. medieps var W. Sm.)

S.

- 187 (mouth of stream "River-wain" near Cenarth OK. 94)  
*Diatoma vulgare*, *Colletonema axinum*,  
*Biddulphia* (hyaline)

- 187? (Scutter bog, Lincolnshire - G.N. 393. 1862  
*Cymbella Ehrenbergi*, *Navicula*, *Utereaphora*,  
*Pinnularia* var??

- 188 (Ponds near Haverfordwest OK. n. 63)  
*Synedra radicans* (var. 2) (in suds)  
 / *Surirella ovata*, small, and large, like *S. crumosa*)  
*Cyclotella speculata*. - (scarce)  
 For medium density - see S. 166  
 For lighter - - - see S. 189

- 189 (Ponds near Haverfordwest. OK. n. 11)  
*Cyclotella speculata* *Synedra radicans*  
*Surirella ovata*, *crumosa* sp., *angusta*.

Lightest parts of S. 188 & S. 166

5.

190 (Mill stream Haverfordwest OK. 55)  
*Syndra pulchella*! *Pinnularia contractum*?  
*aumentatum*!, *Cocconeis*  
*Eunilia arcis* (Scarey)

191 (Haverfordwest OK. 4)  
*Epithemia granulata*!, *Pinnularia oblonga*,  
*Himantidium putinale*, *Cocconeis lanceolata*,  
*Pleurosigma attenuata*

192 (Pond near Haverfordwest OK. 9)  
*Cocconeis Placentula*. *Pinnularia* ~~guttula~~  
*auricularia*? and var? (*P. polygonum* var?) *Stauridium*  
*oblitum* greg.

5.

193 (Hull D. H. munroe)  
*Nitzschia aciculans*, *Cymbella affinis*,

Heavy parts. of S. 359

194 (Bath between Carmarthen & Kidwelly OK. 91)  
~~*navicula elegans*, *pedicellaria*!~~ (scarce) *Pinnularia*  
~~*baileyana* (new sp.)~~ *keegana*, *Mastogloia lanceolata*, *Stauridium lineare*,  
*selina*, *Goscindium minor* (?) *Spirilla crumen*,  
~~*collotrichum eximium*!~~ *Other marine?*

+ see S. 245

195 (Haverfordwest OK. 20)  
*Spirilla pinnata*!, *angusta*!, *ovalis*  
*(eraticula* has been get here) *Pinnularia viridula*!  
*gracilis*. *Cymatoplaera apiculata*!

196<sup>2</sup>. deposit from Lough Mourne Co. Donegal G.M.B.  
 Jun 1865  
*Karula folia*, *obtusata*, *Him. solvirostris* / *planifrons*,  
*Eurost. incisa* w/g (but twice too coarse), *Gomph. subtilis*  
*Fragilaria undata*? (very scarce)

5.

- 196 (church land, fm. N. Haverfordwest OK. 62)  
*Laminariales* (S.V.) *undaria foliacea* (Scov.)  
*Himanthalia elongata* (S.V.) *Nelkia cur-*  
*taula*! (Le S. 17) *Pennularia acrophloea* (Scov.)  
*Stevenson* (*gracilis* Ritter)  
*Lemnaria tetraodon* (Scov.).

- 197 (Same as last, but lighter parts <sup>OK. 25</sup>)  
*Ruppia* is also *Narcissus* (Lambrides), and *N. liliaceus*  
*N. thunbergii* (P.V.) *Luronium bicarpophyllum* (P.V.)! *Him. undulata*  
*area*, *S. C. umbi*, *Pinn. decolorans*, *acrophloea* (Scov.)  
*gitteri*, *Cymbella heterostoma* - *Punt. heterodon* (Scov.)

- 198 (Railway cutting, near Haverfordwest. OK. 6)  
*Cymbella ventricosa*!

5.

- 199 (Gill Bank, Ulverstone, 12:5 h Mif. P. Hodges)  
~~Dactylosporangium sulcigae~~  
~~Lentipes~~ ~~infuscatus~~! (?) *Cymbella maculata*,  
*Pinnularia radiolaria*.

- 200 (Superior W. Smith per OK. n. 80)  
*Rhabdonema minutum*

Widest & deepest part (not  
good now)

- 201 (Neyland, near Haverfordwest OK. 82)  
~~Schizoneura~~ *Grammatophora marina*? *Schizoneura* *gen-*  
*lei*, *cruciger*, *Mitzachia angulata*! *Squadr-*  
*affinis*.

5.

- 202 (Haverfordwest L-Okiden ok. 15°)  
*amphiploea pellucida*.

neatly pure and whole. for drawing  
 on covers & dry mounting.

- 203 (Tidal gathering, Neyland ok. 96° ~~ok. 52°~~)  
*Navicula didyma!* *punctulata*!, Westii?

- 204 (Sussex W. Smith pers ok. 68)  
*Pleurosigma angulatum*!, *Suriella striata*!  
 dead up

5.

- 205 (Pond near Haverfordwest ok. 52)  
*Pinnularia nobilis*, major, viridis (or *Dactylios*)  
*Tellularia*, *Stauroneiforma*, *Odontidium nutibrah*  
*Kutz*, *Suriella biseriate* - *Fragilaria viridis*,  
 - short threads (on the surface)  
 dried up

- 206 (Richmond, Virginia N.W.B.)  
*Coscinodiscus* v.sp. *orthotis*. Suleak

lun 366 - also 635, 636, 637

- 207 (Richmond, Virginia - R. Harrison DMS. 213)  
 nearly same as last but not so clean

among the very weariest was soon of course gone.

5.

208 (Hull Dec<sup>r</sup> 1854 R. Harris) ·  
*Nitzschia sigma*, *Melosira nonmoloides*,  
*Typhloneura nemoralis*

209 (Hull - Octob. 1853 D. Redfern 1854)  
*Typhloneura* sp. var. ? (but *hirsuta*) *Nitzschia*  
*Closterium* (grind) *Cocconeis minor* (very scarce)  
*Pleurosigma fasciata*, *angulata* &  
 209<sup>2</sup>: Brough near Hull - S. Nelson Aug - May 1862  
*Parv. (nav.) varians* (*tenella*) - integra  
 [Hitz. *closterum* the leaves are coarctate]

210 (Ravervordwest 25 Dec<sup>r</sup> 1854 O.K. n. 16)  
*Navicula elegans*! *minutula* <sup>(spectabilis?)</sup> *R. Harris*,  
*pusilla* (then two very scarce) *Pinnularia peregrina*,  
*Hausenii saline*

[rather dirty and too little boiled - does however  
 go well if well toasted in the pan] ·

5.

211 (Ravervordwest 25 Dec<sup>r</sup> 1854 O.K. 140)  
*navicula minutula*!, *elegans* (*scarce*) *Pinnularia*  
*peregrina*, *Hausenii saline*!

[Lighter parts of last ad cleaner, but requires  
 the well toasted Turnip root to get the oil out  
 into the frustules]

212 (Dundee D. Gregory c. 128, 129)  
*Navicula elliptica* Kutz (N. ovalis W. G.)  
*bacillaris* (size 3m.)  
*N. levis* W. G. *angustata*? *denticula tenuis*,  
*obtusa*, *lineata*. *Cocconeis Thwaitesii*, 6pm.  
*tellae* *Scotica*

213 (near Edinburgh Dr. Gregory )  
*Pinnularia pygmaea* Phubbs (id. W. G. - P. name  
 W. G. mol.)

and up I never tried

5.

214 (Deposit from Auvergne A.K.S.)

Dried up &amp; remisted

215 (Kieselgur von Frankenburg, Jaquin per Robert Brown.) - same as S. 78 but not cleaned.

216 / Caermarthen Dk. n. 38)  
Pinnularia viridula Smell.

5.

217 (Schabae Geanw <sup>2d year</sup> J.T. Nunn, )  
Goseinodiscus sp.  
Eupodiumus near Haffsu

218 (From the Kasten deposit, bid y Llo Rheid. Dr. Heming per Dr. Gregory)

219 / Elchies - coarse parts - Dr. Gregory E. 16)  
Pinnularia nobilis t. in, divergens, nodosa  
Tabularia, Scutella, crassata, nobilis,  
splendida, tenax W.G., Stauroneis aculeata  
var. (with parallel sides)

See 157

S.

- 220 (Elektra. Fine parts D. Gregory "E. 16"  
*Navicula coccineiformis*, minute w.g., *Pinnularia*  
*radiosa*, subcapitata. (capitata et) bicapitata. w.g.  
*linearia*, obtusa. *Tabellaria*, *Gomphosphaeria* G. (G. tenella  
 fida W.S. = *Pinnularia aculeata* w.g. non) *Melodus distans* (Litt.  
 in 157) - Stems. ventricosa linearis - acute 3 dentate. Sur. linearis  
 221 (6) ~~dermarthra~~, exposed to tidal influence OK. 26)  
~~Nitzschia~~ ~~harrisi~~? (Scam), *N. linearis* var  
 obtusa? (Sypnium aculeatum Nutz) - vid. S. 333.  
 221<sup>2</sup> (Hickling, Norfolk - F. Kelt)  
*Nav. bacillaris* (Kutz). *Pinnularia cardinalis*, nobilis, *Surirella*  
*erectula* (Kutz) - also *aculeata* (Kutz) and  
*naviculariformis* - vid. S. 333
- 222 (S. Lanes Park - July 55. D. Gregory C. 131)  
*Encyonema prastratum*. *Gomphonema*  
*erectatum*!

5

- 223 (Green Park - July 1855 D. Gregory C. 132)  
 "achanthidium lanceolatum" (not so)

- 224 (Harrowford West T. Okeden - OK. 36)  
*Navicula cryptolephala* var angusta?  
*Pinnularia radiosa* (L.) Blum

at the dinner of 17/8

- 225 (Harrowford West T. Okeden OK. 78)  
*Meridion circulare*, *achanthidium*  
*lanceolatum*

crude. This is for burning on slides  
 and then mounted in balsam

## 5. see end of Book

226 (Near Loch Leden D. Gregory B.52  
*Mastogloia Smithii*, very slender - Navicula  
 angustata?

227 (North base of West Lomond D. Gregory B.57)  
*Siatoma elongatior* (small), *Navicula costata*  
 & *N.G. (Pinnularia integrifrons) Sm.*, *Gomphonema*

228 (St Abb's Head D. Gregory B.59).  
*Epithemia gibba* (very fine) *Gomphonema*,  
 (like *G. punctatum*) *Navicula turbinata*  
 (see 201) *Navicula angustata* blue, *Denticula*  
*ocellata* Sm., *Cyphella testacea*? *Mastogloia*  
*Smithii* very slender

\* for light parts, see 308

5. under a tumulus at Balloch near  
 N. 2/inst in small deposit of Carron fl. - S.P. 1850  
 229 (see 226 & 228) ~~for another deposit~~  
 (see 226 & 228)

for another deposit, for example, see 573

230 (Ulverstone Mrs. Hodgeson /  
*Navicula Lennieri* (front view)! minutula,  
 punctulata! didyma, *Pleurozigma angula*  
 turn *Pinnularia espinus*!, gracilis (seen)  
*Tryblionell. apiculata* <sup>1850</sup> ~~1851~~, marginata, punctata (seen)  
<sup>1850</sup> *Lerma* <sup>1851</sup> (*very rare*), ovata, *Staur. pulchella* (seen)  
 231 (Arctic Sea Lat. 75°-102°, D. Sutherland)  
*Stauroneus pulchella*, *Podost.*

5.

232 (Arctic seas, Assistance Bay, 7 fathoms)  
*Pleurosgma acuminatum*, *elongatum*, *fa-*  
*ciol.*, *pulmatum*? *Pinnularia directa*,  
*Cosinodiscus radiatus*

233 (Arctic seas, assistance Bay, 7 fathoms)  
*Pinnularia directa*, *australis ovalis*

very poor.

234 (Barnickfjæs) R. Kennedy 7:53  
*Ababdonem-adiatum*!, *orthotis marina*!,  
~~mitratus dubius~~ <sup>(3)</sup> *contracta*!, *Navicula mitella*! (var.)  
*helvetica*! ad Kennedy, *buccinaria diafragma*,  
*Pinnularia distans*!, *campylodes-*  
*eus Hodgsoni* (very scarce), *Bidulphia rhombus* (very  
scarce))

5.

235 / *bumbræ* R. Kennedy /  
*Goscinodiscus radiatus*, *concinnus*, *O. pedice-*  
*Ralfsii*! a adsp WG., *craspis*; *Ababdonem-adi-*  
*atum*!, *Navicula* <sup>Humerosa</sup> ~~angulosa~~ <sup>WG?</sup>, *Smithi* (*ovalis*)  
*Sm.* (var?), *Lyrat* (~~small~~ <sup>a. b. 8 B. 6 Spec.</sup>), Kennedyi (small Corp)  
*Stauroneis pulchella* <sup>B</sup> (common)!

236 / *bumbræ* - very light parts of flesh)

237 (Barnickfjæs) R. Kennedy 7:53  
nearly the same as S. 234, but less *Navicula*  
and *nebulosa* <sup>(3)</sup>. *Tugonilla dubia* <sup>(2)</sup> *contracta*, *Coccone-*  
*major*, *aperta* <sup>purple</sup> (*allegheny*), *orthotis marina*, *Aba-*  
*denem-adiaticum*, *grammatophora* *lesper-*  
*tina*, <sup>subtilissima</sup> *marina*, *gen. Eupodiscus* *sulcifrons* (new)  
*Flavos. formosa* var *angusta* (*decorum*?)

S.

238 (Ulverstone Mif E. Hodgson No. 51)  
*Navicula cryptcephala, Pinnularia radiosa,  
 acanthodium lanceolatum*

5

241 (Isle of Man Mif E. Hodgson No. 28)  
*Synedra Gailloni*

239 (Hillfoot, Cumbria 5:54. A. Kennedy)  
*Acanthodium lanceolatum !!*

242 (Hayland shore of Miford Haven, F.O. Kuden Oct. 79)  
*Biddulphia aurita*

240 (Ulverstone Mif E. Hodgson No. 61)  
*Synedra affinis, Biddulphi-aurita var.  
 (Eupodium radiatus, extremely scarce)*

243 (Mill stream near Haverfordwest. F.O. Kuden Oct. 71)  
*Ceratia circus, Melidion constrictum var.,  
 Surirella amphioxys (as in S. 244), Beyebell  
 ventricos-*

5.

- 244 (Spring near Haverfordwest, T. Okeden. OK. n. 48, 49)  
*Pinnularia*<sup>+</sup> [*lond. constricta*] *Surirella*  
*amphioxys* W. Sm. ! *Cymbella ventricosa*

+ same as S. 295

- 245 (Wall within the town of Haverfordwest OK. 33)  
*Navicula pusilla* (fide Sm.)! *Achnanthidium*  
*coactatum*! *orthotrichum mirabile*! (Plutejul)

see S. 513

+ see S. 194

- 246 (Ulverstone, Gillbank - Mif P. Hodgson no. 21)  
*Meridion Zenkeri*, circulum 8<sup>o</sup>

5.

- 247 (Wall within Haverfordwest T. Okeden. OK. 34)  
*Navicula pusilla*! *Achnanthidium coacta-*  
*tum*! *orthotrichum mirabile* (very scarce)

obtained by burning the material of S. 245

- 248 (oysters from Milford Haven, T. Okeden OK. 77)  
*Eupidiaceus erafae*! *Coscinodiscus* - - -

- 249 (Garmyle 6:55 R. Kennedy)  
*Gomphonema curvatum*! *Eurotia ericetorum*!  
*(Surirella eraticula* small and very scarce)

5.

250 [Lightburn 3 m. S. of Glasgow 6:55 A. Hennings]  
*Diatoma vulgaris* good and pure

251 [Dugdale Cork Harbour 12:7:55 W.S.]  
*Syndra (Totaria) undulata*, Nitzeckien  
*Specularia amphiflora*. Ligmoidea

\* 100 staves from (about 50) broken up in & other larger  
 pieces - Balsam

252 [Barley Lough, Co. Cork 4:7:55 W.S.]  
*Goniphonema rostratum*, *elongatum*, *dictyon*  
*grandis*, *lunaria*, *diadema* ad *bidentata* (lost  
 very scarce). *Syndra ulna*?

5.

253 [Lough Avail, near Glen Gariff, Co. Cork - W.S.]  
*Goniphonema subtile* Schle. *Epithemia rupestris*,  
 ad *E. tetraodon* ad *diadema* (lost scarce)  
*caribella membrana*

254 [Cattlecary, among Sphagnum 23:6:55]  
*Pinnularia gracillima* W.G. *Epithemia rupestris*,  
 lost, few

255 [West Cork, off ~~Shadee Island~~, off Sargasum Dr. Bailey]  
*Syndra (Totaria) undulatum* -

5.

256 (Deposit from Willington, Connecticut, Dr Bailey)

257 (Duval's creek in Lake Monroe, at Fort Enterprise, Florida  
~~excluded~~ Serpinae mucosa, Noricul. ? (Stauroneis macula-  
 tor Bailey)

<sup>2</sup> 257 (Lucernmore near Lancaster C.J. March 56)  
 Pinnularia stauroneis formis var. ??

258 (Carmyle in old Quarry 7:55)  
 Gomphoxema contractum, acuminatum,  
 Sphaeridium radicans, leucostoma var.

5.

259 (Minehead, Somerset, May 1. Gifford 8:55)  
 Navicula (like N. angustata) &  
 Pleurosigma attenuatum, Encyonema prostra-  
 tum (small) Pinnularia radiosa, gracilis, &

260 (Bed of the Tummill - D. Greville 17 coarse)  
 Gomphoxema germinatum, Tetracladus emer-  
 ginated!

261 (Bed of the Tummill D. Greville 17 medium)  
 Tetracladus marginatus, Achmatthes exili,  
 Gomphoxema germinatum (few) Cyathella  
 truncata W.G.?

S.

- 262 (Bed of the Tummell at Giville No 47, light)  
*Achnanthus apiculatus*, *Tetraeckia emarginata*,  
*Symbella hemisphaerica* Wg?

- 263 (Yen Fang 1:9:55)  
*odontothrix medior* i., *Sympetrum ulm*

- 264 (Kinroft, III near the Lake 29:8:55)  
*Fragilaria capucina*.

For crude see 260

S.

- 265 (Kinroft, No V near the Lake 28:8:55)  
*Navicula cuspidata*, *Pleurozigma laevistriatum*?

- 266 (Finnity near Lock Leven No II. Heavy 27:8:55)  
*Epithemia gibbe* var? (See S. 274) var  
*parallela* Grunow

see S. 274

- 267 (Finnity near Lock Leven No II. Light 27:8:55)  
*Symbella* . *Coccococcus cyathiformis*  
*Macrocystis*

in S. 274 for light Relv

S.

- 268 (Finnity near Lar Lough n<sup>o</sup> W 27.8.55)  
Naricula Cymbella

crude (see 258)

- 269 (Roa Island near Silverstone Miss P. Hodgson)  
Grammatophorinae!

- 270 (Roa Island, near Silverstone - off Clevedon)  
Dorothyamphiceras

very dirty

S.

- 271 (Small pond in Glen Selt at Shooting Lodge - See 53)  
Epithemia hydnomanni, proboscidea (Seary)

Heaviside's

See 216 and 302

- 272 / Fell End. per Miss P. Hodgson  
Himantidium pectinale var. adnatum meso,  
Tabularia flocculosa

may be mounted in Balsam, if well  
soaked in Turpentine

- 273 (Silverdale, Lancashire - off shells C.T.)  
Pinnularia borealis, Nitrodothia amphioxys!  
Cymbella testicula var obtusa, Mastogloia  
gerillae! Epithemia gibba var

5.

- 274 (Tinmity near Leek Lever O' Heavy past 27:8:55)  
*Epithemia gibba* var. *Cymbella scitica*, <sup>*helvatica*</sup>  
*-conema cyathiforme*, *Navicula* <sup>*substrata*</sup>  
 see 5. 266

- 275 (Hunthill near Elches, in a spring D. Gregory E. 14)  
*Pinnularia modesta*! *nobilis*, *Dactylios* (Scare) *major*  
*anceps* ? W.G., *Surirella nobilis* (Scare) *biseriata*,  
*Gomphoneme capitatum* ? Sm

medium and lighter parts of nest.

- ~~276 (Blackhead deposit coarse parts D. Dickie)~~  
*Cymbella Ehrenbergii*, *Epithemia longidens*,  
*Hemicula* ~~conema~~ <sup>(Scare)</sup>, *Surirella nobilis* (Biserial)  
~~varicosa~~ <sup>large</sup> ~~now~~

- 276 (Hunthill near Elches, in a Spring D. Gregory E. 14)  
*Pinnularia nobilis*! *Dactylios*! *major*, *Surirella nobilis*,  
*biseriata*

coarse parts and requiring the air pump.

5.

- 277 (Lincolnshire <sup>5th H. munroe</sup>)  
*Navicula tenuissima* var? (*nostauros*), *Cyclo-*  
*tella speculata*, *Typhlonella angustata*! (small)  
*Denticula tenuis*, *Cymbella helvetica*, *Gomphoneme*  
*capitatum* ?

- 278 (Birdhope, Redesdale Northumb<sup>r</sup> D. Guille <sup>65</sup>/<sub>86</sub> Sept 55)  
*Cymbella regularis* ? (like *Nav. angustata*), *Nav.*  
*cule* ~~angustata~~ <sup>of</sup> ~~an Cymb?~~ *Mastogloia*, *Smithia*  
*odontidea* *Harringtoni*, *Epithemia* v. sp. *Spelaea*  
*argus*? (or *vallata*, or *mish*) - *Denticula* n. sp. ?

- 279 / Peterhead deposit, coarse parts, D. Dickie,  
*Cymbella Ehrenbergii*, *Epithemia longidens* &  
*stabilis* (smthg. irregular & wavy) <sup>and</sup>  
*Navicula* ~~semene~~ <sup>large</sup> ~~Nav. Semene~~ <sup>large</sup> ~~Sm. parts~~  
*Pinnularia* varian W.G. fig. 28) - Littoral.

S.

280 (Peterhead Deposit - Fens & pools D. Dickie)  
*Epithemia turgida* *cyclotellus* *antique* !, *Navicula*  
*semenv. Sm. part.* / = *Pinn. varium* W.G) *Odontidium*  
*multilete*, *Tabellaria*, *Navicula laevisima*

281 (KircKnacullen, Co. Cork Prof Smith)  
*Navicula amphibiaene* !

requires air pump, or the well soaked in turpentine

282 (Peat moss, Ulverstone M. J. Hodgson)  
*Spirorbis amphioxys* ! ex

S.

283 (North mountain, Nova Scotia Heavy peat. OK.)  
*Stauroneis Baileyana* ? *Pinnularia notabilis* &  
*Nav. stabilis* (*amphib. navicularis* Ehr.) !

284 (North mountain, Nova Scotia, medium density OK.)  
*Stauroneis Baileyana* ? (small), *Navicula semenv.*  
*naviculae* R. & A. (N. *lema* Sm. partim -  
*amphib. navicularis* Ehr. !)

285 (North mountain, Nova Scotia, Light peat. OK.)  
*Navicula multicaulis* ! (small with a false  
*staurus*)

S.

286 (Haverfordwest. T.Okeden 4:2:56 OK. 28)  
*Nitzschia amphioxys!* *Pinnularia haunaei*.  
*famili!* *Surielle angustata*

287 (Milldam, Haverfordwest. T.Okeden 4:2:56 - OK. 18)  
 - *Surielle* *O!* *S. amphioxys!*, *splendida!*, *cocconeis* *placentula* (strue distinct,

Light parts of S. 183

288 (Bog near Caermarthen T.Okeden 4:2:56 - OK. 101)  
*Gomphonema tenellum*, *Syndra fasciculata*??

S.

289 (Haverfordwest. T.Okeden 4:2:56 - OK. 86)  
*Fragilaria revoluta*!

much water for sake of the S. veins

290 (Haverfordwest T.Okeden 4:2:56 OK. 84  
*Surielle pinnata!* *panduriformis!* *achmauthidium*  
*lanceolatum*, *Pinnularia mesolepta* (scarce)

291 (Lüneburg Deposit. mis. E. Wallsgroen)  
*Cyclotella minutula* *Corbicella rhombifera*  
*orthosira arenaria*, L.

Differ from 311 by the great want of *Syndra*

5.

- 292/ near Ulverstone Miss E. Hodgson  
*Mastogloia lanceolate* fide Wdm (auf Pl. M. Dendri)  
*Pinnularia* *Cypria*, *pequena*, *Achnanthus* sub.  
*septaria*

- 293/ Exmouth Miss Bellings  
*Diatoma vibuum* : Lomophora  
*flabellata*

crude

- 294/ Haverfordwest T. Okeden OK. 105)  
*Pinnularia radiosa* !  
 stipe rather too close for sl: an Collat. neglect?

in S. 369

5.

- 295/ Neyland, N. Haverfordwest, within reach of the tide  
 T. Okeden - OK. 106)  
*Pinnularia* ! (near *O. radiosa* but  
 broader and more coarsely striated) same as in S. 214

- 296/ Baermarken bog T. Okeden OK. 108)  
*Stauroneis gracilis*, *bimella* *affinis* *emarginata*  
 up to Nag.

- 297/ Neyland N. Haverfordwest T. Okeden. OK. 109)  
*Seriella angusti. ovata* (small), *Nitzschia*  
*Palea* !

See 390 and 391

S.

298 / Bog near Baerenthal F.Okeden OK. 114 )  
*Navicula cocconeiformis?* <sup>*varifima?*</sup> *bacillina?*, *rhomboides*  
not seen  
*(large), evaginaria?* (*not constricted and capitate ends*)  
*Gymbella affinis* - *Pinnularia*: several, *Stauroneis*  
*aniceps* &c

299 / Haverfordwest F.Okeden OK. 120 )  
*Synedra ulna?* *fasciculata*, *Colletome vul-*  
*-gare* (*frustulus*), *Matoma elongatum* (small)

for crude see 353

300 / Bog near Baerenthal F.Okeden OK. 112.B )  
*Navicula* <sup>*affinis* Sm.?</sup> *formosa* var., *Luriella tenera* W.G.,  
*Meridion constrictum*, *Fragilaria viscens*

5.

301 / Bog near Baerenthal F.Okeden <sup>*light*</sup> OK. 112.4 )  
*Stauroneis* <sup>*variegata*</sup> *variegata* with lines on the last  
*Pinnularia* (*intercepta*)

302 / Haverfordwest F.Okeden OK. 159 )  
*Nitzschia linearis* !

crude of rest.

303 / Haverfordwest F.Okeden OK. 159 /  
*Nitzschia linearis*

buried, fm below

5.

304 (Haverfordwest T. Okeden  
*Gomphoxeme germinatum*!) OK (?) hauld

305 (Haverfordwest T. Okeden  
*Synedra pulchella*!) OK, non kept

end of next

306 (Haverfordwest T. Okeden  
*Synedra pulchella*!) 5th, non kept.

Same as last but prepared for balsam. See 302  
 OK. Did not keep away, as he had plenty  
 within OK. 55 = ab. S. 190

5.

307 (Jersey <sup>Mf. Beaufort</sup>)  
*Grammatophora marina*, <sup>\*</sup> *Synedra Gailloni*!

\* This form has fine striae, see S. 339

308 (Victoria Bank, Kew - lightest parts, Feb 1856)  
*Stauronitis* ? (minute), *Dicadesmus*  
*peregrina* W.M. (Side view)

For heavier see 356

309 (Chabot Guano 1856) light parts  
*Synedra* ? (partly broken & whole)

(not much) (after broken & whole lost  
 it must be the rest, for tab. 309  
 is based on the top)

Separated from 375 to show the Synedra better -  
 the put on the slide with a glass rod

S

310 (Swarthmoor, Ulverston - Miss E. Hodges)  
*Surirella crenulata*, *Cymbella maculata*,  
*Syndes. ulna*, *Cocconeis cistula*.

312 deriges 2. Okto. ok. 185<sup>h</sup>  
*Rudeusinae*, *Lamprolium*, *Cymbella Strumigenys*

311 (Ulverston - Brackish water - Miss E. Hodges)  
*Surirella crenulata* ?

312 (Pier on Roa Island, N. Ulverston Miss E. Hodges)  
*Campylodiscus parvulus*, *Nitzschia sigma*

S.

313 (Deposit from Old Man Muntan, Lancashire Miss E. Hodges)  
*Eunotia diademata* - *Himanthidium lideum* & -

314 (Haverfordwest 7. October x  
*Cymbella scutella*

315 (Haverfordwest 7. October (slightly lost))  
*Himanthidium festucale* ?

same as 351 (for crude see 349)

S

316 (Haverfordwest, F. Okeden OK. 142).  
*Synedra tenera?*

Crede (for bild. see S. 331)

317 Haverfordwest. F. Okeden OK. 146)  
*Nitzschia acicularis* (Pall.) Mittw. var.  
 Dried up & remastered, looks wrinkled

on fol. N. acicularis var. erosa (nabamca 3902),  
Crede

318 (Haverfordwest. F. Okeden OK. 131)  
*Nitzschia* (<sup>Diplorella</sup> <sup>common var?</sup>)  
 (Cane, oblique var. Nutt. (in v. f.))  
 (Syn. <sup>subtilis</sup> Schleiden var. lineariflora Steyermark, 2)  
To be well soaked in tap water to get  
out the air from Pinn. midrib &

S

319 (Haverfordwest. F. Okeden OK. 133)  
*Cymbella ventricosa* very minute

found with 319 (OK. 135) -

320 (Neyleand near Awest F. Okeden OK. 149)  
 (synonyme v. <sup>longula</sup> <sup>longula</sup>)  
~~Nitzschia longula~~ <sup>longula</sup> or *N. tenuissima*  
 The parasite is very cavous on the back, and flat on  
 nearly so in the lower side. It is white like *N. tenuissima*  
 or *N. tenera*, but much smaller. It is when an adult - a  
<sup>cooper</sup> <sup>of</sup> <sup>an</sup> <sup>old</sup> <sup>specimen</sup>  
<sup>adult</sup> <sup>it</sup> <sup>is</sup> <sup>reversible</sup> see S. 384 Crede (for bild see next)

321 (Neyleand near Awest F. Okeden OK. 150)  
 same as last, but bristly

S.

- 322 / Haverfordwest - T. Okeden. F.O.  
*Cymbella ventricosa* T. V. - occident? navicula  
*Cymb. Helvetica* - - navicula? minuta

Crude / Varibiles S. 323, S. 32 //

- 323 (Haverfordwest T. Okeden - F.O. 157)  
*Cymbella ventricosa*, *Helvetica*, &c.

- 324 (Neijland near New T. Okeden - F.O. 140)  
*Tayblionell. lindenii* W. Sm. (since as  
 caudate, but slightly more numerous than in Smith,  
 show)

Found with 384, 385 and 386

S.

- 325 (Neijland near New T. Okeden - OK. 152, 153)  
longer and inflected  
*Denticula* stulta ad *Cymbella maculata*?  
very minute

- 326 (Swarthmoor, Ulverstone mp. E. Hodgs.)  
*Nitzschia linearis* B. N. S. (n. mod. *N. Sm. olens*)  
*amphifleur-pellucida*, *navicula* *tumid*

much mixed

- 327 (Haverfordwest T. Okeden - OK. 158)  
*Navicula* ?? (very minute)

very light parts of S. 323.

S.

328 (Deposit from ~~Holyhead~~<sup>Holyhead Lake</sup>, Caithness, S. Peacock)

buccinum Thwaitesii and var with an intense  
square nodulus Nav. macul. W. G. ?, Navicula pustula,  
obtusa! Denticula obtusa & brown <sup>guttular</sup> <sup>argus</sup> <sup>proboscidea!</sup>  
Smithii, Epithemis ocellata, Zebra, ~~granata~~ & v.

+ P (first heavier density only see S. 328) <sup>(Hornbeam shell)</sup>

329 (Bologas, Penzance. J. Alfs )

Fragilaria variegata with Himationium  
spectabile minutum.

330 (Hoe moor, Penzance - J. Alfs )

Himationium spectabile, all forms and sizes!

+

S.

331 (Haverfordwest. T. Okeden. OK. 142)

Synedra tenera?

(For crude see S. 316)

332 (Well at Papsage, Co Cork. Prof. W. Sm. R.)

"Fragilaria capucina" prof. W. Sm. - al  
potius Fr. variegata var.

333 (Swansea L. Okeden OK. 102)

Mitrochis unicarinata obtusa? (seen  
also Synedra acuta of Kutz.) - also  
mit. amphioxys, Tryblionell <sup>near</sup> minima?  
Amphora marina? acanthid. coriacea

see S. 221

fresh water

5.

- 334 (Bog near Penzance J. Ralp,) *Synodre biceps* T. V. - also var? (Eunotus - flexuosus Kutz. Scare) Himant. major (Stein)  
arcu. und.

nearly same as next, but not so clean  
and with more of *Syn. biceps*

- 335 (Bog near Penzance 17.5.56. J. Ralp,) *Synodre biceps*, and var? (Eunot. flexuosus Kutz)  
Himant. major, arcu., undulatum.

5.

- 337 (Dolgelly, Wales. J. Ralp) /  
*Tetraclitella lacustris* - Himant. undulata

Crude (for boiled see next)

- 338 (Dolgelly - Wales J. Ralp) )  
same as last, but boiled.

- 336 (Caermarthen T. Okeden OK. 161) *Cymbella cuspidata* very fine & firm -  
striae on the dorsal half only 20 in 001, on the ventral,  
about 24 or 25. - So intermediate between *C. granularis*  
and *C. cuspidata* Sm.

- 339 ( Neyland, off Rock - T. Okeden - OK. 172) /  
*Cymbella maculata* of Sm. the figure, but  
not of this slide

S.

- neyland near  
 340 (Haverfordwest F. Okeden ok. 276)  
*Cymbella vertic? macul?* (same as in 5.325)  
Small

S.

- 343 Caermarthen - F. Okeden - ok. 24)  
*Nitzschea amphioxys* 3 much smaller than  
 usual ("*N. minutissimum*" F. Ok.). also *N. amphioxys*  
true

- 341 (Penzance J. Alfs.)  
*Himantidium majus*, *Synidr. bicuspis*

- 344 (Haverfordwest - F. Okeden - ok. 182)  
*Synidr. minutissimum*

Burnt and washed below  
 F. Vines.

- 342 (Haverfordwest F. Okeden ok. 179)  
*Synidr. longissima*, *buxeratiformis*  
~~gigantea~~? *Himantidium arcum* (Heyford)

- 345 (Kingshorn, Fifeshire (fresh water) D. Reville 7:53)  
*Synidr. ulna?*

S.

346 (Utracornia - S. Ralp.)  
 "Amphifleur scalaris" for S. Ralp.  
 but Conn not so, but only A. rufescens & others

S.

349 (Haverfordwest I. Okedur Oct. 27/21)  
 Synod. tenella? and radicans

crude.

347 (Cumbrae 30:5:56 R.H.)  
 Schizoneura Smithii

(for crude see 436)

348 (Cumbrae 30:5:56 R.H.)  
 Schizoneura Smithii

(for crude, see 437)

350 (deposit from Old man Manton - Miss. Hodges)  
 Surinam bilobata, &c.

coarse parts - very diff. from S. 313

351 (Muskland Canal May 1858.)  
 Diatoms vulgaris pure

5.

352 (Haverfordwest salt marsh I.Old. ok. 176)  
*Nitzschia scalaris*? (leaves - 2 or 3 mm  
 wide) *N. obtusa* (very scarce) *Navicula pusilla*  
*elegant*, *Pinnularia peregrina*!, *Pleurozium*  
*obscureum* (only seen & named) -

353 (Caermarthen F.Olden ok. 177)  
*Cymbella cuspidata* ann. ok. 161 (S. 336), but  
 not so pure, also *Spiroderma linearif.* and *Staur-*  
*omopsis var. fragil. viridis*

354 (Caermarthen F.Olden ok. 178)  
 Light parts of the last

5.

355 (Haverfordwest 2.Olden Penwith (M))  
~~*Achnanthus gracilis* H. & A. var.~~  
*Marsilea quadrifolia* (L.) 5<sup>2</sup> and 6<sup>2</sup> - *Fragilaria*  
 ann. 388, *Denticula*? *petalophora* - see 245  
 for crude see 445

356 (Penzance I. Nalff June 1856)  
 " *Amphipleura inflexa* Bréb." - ~~conf.~~  
~~*Pleurozium anomolus* Schreb.~~  
 since seen with a  $\frac{1}{2}$   
 (for listed from Wales see S. 472)  
 for crude see 466

357 (Lancaster - salt pools C.J. June 56)  
*Colletotrichum acuminatum* var.

357<sup>2</sup> Stallingborough near Great Gurney, Lincoln  
 Colletotrichum <sup>G. Norm.</sup> 362<sup>2</sup> specimen ~~W.M.S.~~ - Body of  
 108<sup>2</sup> tied up & remained - very  
 bad & dirty - may be broken away

S

- 358 (Neyland F.O. Keden - OK. 171)  
*Susirella* sp. ~~*lineare*~~ ~~*lanceolata*~~ ~~*lanceolata*~~ Hull. Pinnular  
 hemiptera? - *Himanthidium arvense*  
 (below difficult to enter)

- 359 (Hull D.H. murrayi)  
*Cymbella* aff. ~~*affinis*~~, *Hitzschia linearis*  
*Synecr. radians*  
 see 5.193 (for key pls.)

- 360 (Hull D.H. murrayi)  
*amphibalanus* daniellii sp. (or *Schizoneura*  
 sp.) same as in 115 and 3.41

+ also from G. Normar 58, & mixed

S.

- 361 (Kewerfordwick, Salt marshes F.O. 195<sup>1</sup>)  
 slightly separation from 352 in the crude that  
*Thuretia obscurum*. *Cyathella kutzingeri*

- 362 (arran. Broadick Bay - aut. 53)  
 .... *Navicula Lyra* (very scarce) - also *N. fimbria*  
*N. praecincta* W.G (very scarce)  
 heavy parts of a colony of *Schiz. crinoidum*

- 363 (arran. Broadick Bay off Lunderley aut. 53)  
*Homocladia* ( ) mixed with  
*Schizoneura* *Dillwynii* ( )  
 also some pieces like of *Cheiloceras* W. Hammar

light parts of last.  
 called by Prof. Dr. "Sch. implicata?" and by me  
 formerly *S. crinoidum*

5

364 (Springhead near Hull - D. H. Murray)  
*Synedra ulna* sp? meridionalis

365 (Spring head, near Hull D. Murray)  
*Synedra delicatissima* <sup>*radiata* sp?</sup>, *Cyclotella*  
*minutula* Kutz [very indistinctly marked]  
 and punctata

~~Aug 20~~

366 (Well near Douglas, J. Y. Mac. Miff. Hoelzer)  
*Fragilaria varia* var "F. capucina" sp? <sup>W. S.</sup>  
*leptostoma* sp? <sup>Hoelzer</sup>

5.

367 / Cenularia  
*Schizogonium dillwynii*

(for crude see 422)

368 / Aberdeen D. Murray  
*Schizogonium dillwynii*

- boiled for 45 min some on 423, or  
 for crude see 426.

369 / Haverfordwest T. Okey OK. 201  
~~Pinnularia radiolaria~~ ~~Collotrichum angustatum~~  
 Pinnularia radiolaria var.  
 thick fine - surely must be ~~Collotrichum angustatum~~

see S. 296

S.

370 (Hull

*Pinnularcia aerocephala*? Nitz. curvula s.n.  
 (Syn. *nitens* (L.) L.)  
*Neurosigma attenuata*

D. Murray

371 (Gourock, on leaves of Zoster, 24:41. 5'2)  
*Schizoneura helminthoides* decolorans

(holotype from dried specimen)

372 (Exmouth June 1886 Miss Cutler)  
*Schizoneura dillwynii*

372<sup>2</sup> Watermouth Cavern, off Exmouth, exuv. J.S. 2:65  
*amphipora confusa* (entom. frustulus - *Dimerogramma*  
 (Derm.) *arcuata* n. sp. (J. Staunton) - Nitz. venosa n. sp. (frustuli  
 amphip. ligata) (exuviae see 430) - *amphip.*  
*riphaea* (in very light)

S.

373 (Haverfordwest T. Okeden - F.O. 197)  
*Kiamantidium pectinale* large

(exuviae see 440)

374 (Stream at Toward Point 27 June 1856)  
~~at the~~, near ~~at the~~ mouth  
*Synedra* gadana meridionalis circula. 8  
 varicula tumida &c.

375 (Pembrey - Miss Cutler 1:7:56 - I. 1)  
*Schizoneura Smithii* - frustula long.  
 also *S. molle* ? (smaller broader, and closer  
 striae!)

(exuviae. see 447)

5

376 (Exmouth - Miss Cutts F. Linn 1858)  
*Schizoneura Smithii* ordinary sized punct  
 Lent as *S. torquata*  
 (for crude see 448)

377 (Exmouth Miss Cutts G. Linn 1858)  
*Schizoneura molle*.  
 Lent as *S. Wyattice*.  
 (for crude see 449)

378 (Pond at Haverfordwest F. Okeden. OK. 240)  
*Stauromenis dubia* W.G. (*Navicula decipiens*,  
 OK.) is a species of *Achna* ?? (no stauron to upper & lower)

5

379 (Pond at Haverfordwest F. Okeden 240  
*Stauromenis dubia* W.G. lent three over  
 the strainer - an Navicula sp. *N. decipiens* sp.  
 379<sup>2</sup> Wall of W. Norman's Barn Hull - solid and cloudy  
 of 15.495. 9.V.12  
*Nav. appendiculata* Kutz - ~~Griffith~~ *Normanum* Rob  
 (Comb. arm. W.G.) same as in 3.67

380 (Silverdale, Lancashire C.T. 18.7.56)  
*Synedra ulvae* var  
~~ulvae~~ ? - wider linear; pointed  
 short - ends as in *S. ulvae*, and no distinct radial  
 vacuole in the middle

381 (Well at Haverfordwest, F. Okeden. OK. 232  
*Himantidium major*

(for crude see 455)

381<sup>a</sup> was a small quantity never looked by Mr  
 Okeden (his OK 232), but all mixed with 3381 (stone-like)

5.

382 (S. side of Grinnan Canal between Ardrosshaig &  
Lockfield, gently washing bottom. mychum 16:7:56)  
Diatomell. Balfouriana!! Epihemia alpestris, superstis!  
ocellata, gibba, Denticula tincta! Coc. Thwaitesianus!  
Gymnophenia vibrio!, Cyclidella sativa var. obtusa & helvetica

383 (Coarse density of glass) 16:7:56  
Epihemia alpestris, superstis, ocellata, gibba.  
P. alpestris seen. G. libula Ehren.  
swamp & remnant land not good

384 (Neyland T. Okeden OK 237) n.  
Alphidion peradice, Stratella  
unipunctata.

crude - for burning in com  
not very good

5.

385 (Neyland T. Okeden OK 267)  
Stauroneus saline! Navicula lessonii!  
Pinnularia cyprina

386 (Neyland T. Okeden OK 268)  
Pleurosigma angulatum var? Scandigaster  
(P. estuarium var?)

rather much broken - see

387 (Fell End C. J. July 1858)  
Navicula mutica Kutz?

among oscillatoriaceae in a trough

5.

388 (Salthouse near Ulverstone Miss P. Hodge)

*Nitzschia Palea*? or a spec of *Homoecidium*. .)!*Nitz. dulcis*! var<sup>\*</sup> (resembling *Typh. acuminata*)*Susirella v. sp.* - *Synedra minuta* - &

\* more stratus; same as in S. 103, S. 457, 296, and 520-2

a great mixture of salt &amp; f.W. species.

389 (Haworthdown L. Oxford 100 m.)

ad nat. dim. mesodon

390 (Salthouse near Ulverstone. Miss P. Hodge)

*Odontidium mesodon*

5.

deposit from

391 ("Linton in Prussia." D. H. Munroe)

*Cyclot. rotulae?**Olkoszia arenariae*, *Cyclotella punctulata*.*Cymbella strenua* Fr.

I suspect this is from Dafsan Kielder,

and probably Eng. tab XIII. II. #1

392 (Rocky ground S. of Arrochar 2: 8156)

*Cymbella effigie*, *Helvetica*, *affinis*, *Coccone**Ihwaiteana*, *Gomphonema dichotoma*, *gemellum*,*Himantidium arcuatum* Fr.

393 (near Ambleside, Lake District, 1 m. by Bowes Mr. Batty Bury)

*Synedra tenera*

for crude in L56

5

394/Corrie hills, Aran- 7:56 (R. Kennedy)  
Odontidium hymenoides

see 186

395 New Lancaster C.J. auf 56)  
~~Nitzschia lanceolata (Mull.) St. Sigma.~~  
~~N. dubia / var. lsm (Tryblionella apiculata n.sp.~~  
Same as in 319)

395?/Pools under Capstone Hill. Ifromby p. 186<sup>4</sup>  
Nitzschia heterostoma Grunow (as var viscidum N. heterostoma)

396/Lock Vack, Hill Krauter 1400ft R.R. A.K.G. 2114)  
Cyclotell-<sup>opuntia var</sup>! Stauroneis pulchella?  
Synedra tenem. (coccoeithwaitesii / and var  
with a green stain var. maculata?). Cyathell  
maculata Epithemia fasciata Sorex, turgida

+2

5.

397/Corrie hills, Wash. from 2000ft Crail. A.K.G.)  
Synedra gallica var. S. gallica (var. 719)  
Stauroneis pulchella?  
and <sup>leptostoma</sup> ~~leptostoma~~? Eupodiscus ralfsii,  
Surirella fastuosa -

398/Hills S. of Lock Scummell <sup>2000ft elev</sup> A.K.G. 247 1.  
Pinnularia. Stauroneiformis var., lata (scum)  
Acianthidium gracile (?)

399/Seaton, S. Devon. <sup>Antiup Hodge</sup>  
Stauroneis pulchella? Synedra gallica. Gramma  
-lophora serpentina. marina <sup>\*</sup>? Rhizidium mearensis  
(also but extremely rare Campyl. parvula, orthotrichum  
marina, Podostem. montagnei & maculata. new  
cyathella undulata

\* There are two forms here: one has twice the air-sacs,  
about 3 mm <sup>long</sup> <sup>wide</sup> 0.01).  
by Noctis N 3: the other much finer, and regular the first <sup>+</sup>  
but the last is not so difficult as in S. maculata.  
and is same as in S. 539

5.

400 (Lizard Point

J. Ralff).

*Epi themis longicornis, argus, turgidae, pro.  
bisidea, !!**Mastagria Granular. cornuta*

401 (Lizard Point

J. Ralff)

*Pleuralgia transversa, Empodium exasperatum  
Symbionellus contractum, Cocinodiscus radiatus  
variable Br., Triceratium attenuatum (very rare  
orthoseta murina, Symbionellus contractus sensu), Codium  
maculata*

402 (Collingham near Hull

D. Mummae)

*Pleuralgia attenuata*402<sup>2</sup> (Shapney near Great Malvern Station, Buells. Ok. 560)  
*Pleuralgia attenuata sp., Coccocon, Placunula (and  
and sporangial) Ament. fl. elliptic & lobed var. gibbera  
for dry moisture**some marked S-402<sup>nd</sup> among the heavy parts of the  
gathering.*

5.

Aug 1856

403 (Hawerfordwest T. Okd, Ok )

*Narcissus diephile :. Romanthus pechi  
al? (small)*

404 (Banff - Ba-1. 12:8:56)

*Siatoma vulgare! Coconotus Pediculus! Gum.  
-phonema contractum, curvatum**no. 2. . . . .*

405 (Pool S. of Aberdeen. A-1. b. 14:8:56)

*Melotira subfusilis, Epi themis granulata,  
Synedra acicularis, gracilis, &  
Fragilaria rupestris.**for crude no 457*

S.

406 (Pool S. of Aberdeen A. 2. 14:8:56)  
*Melosira subflexilis*, *Lepid. aciculare*  
*gracilli*, *Epithemia granulata*, *argus*, &c

same pool as 5. 405

sometimes ~~so few very long frustules~~ (*perangul*) <sup>spitt</sup> ~~green~~  
*anellae*

407 (Rock S. of Cove N. of Aberdeen A. 3. 14:8:56)  
*Denticula obtusa*, <sup>(brown)</sup> *Cocconeis circula*, <sup>(brown)</sup> *Navi-*  
*-cula elliptica* Kutz. (ovalis Sm.) <sup>(brown)</sup> *Cocconeis Swartzii*

nearly same as 39. See also 4. 480

408 (Speling S. of Aberdeen A. 5. 6. 14:8:56)  
*Orthodesma orichalcea*, *odontidea mediana*  
*mutabilis*, <sup>(brown)</sup> *Fragilaria virescens*? (like adm.  
 mutab. but thick walled), *Diatoms elongata*  
 var. ?, *Achnanthidium lanceolatum*

found in 458

S.

409 (Cave at Cove N. of Aberdeen A. 6. 1. 14:8:56)  
*orthodesma Dickii*, ~~odontidea medialis~~  
*Fragilaria virescens*? *Himantidium*  $\approx$   
*Achnanthidium microcephalum* (Scen.)

very tightly built, & require blue well coated in tufts

410 (Cave at Cove N. of Aberdeen A. 6. m. 14:8:56)  
*orthodesma Dickii*, ~~odontidea medialis~~ *Fragilaria*  
*virescens*? *Himantidium*  $\approx$   
*Achnanthidium microcephalum*, very seen

1st much build - heavier parts

411 (Cave at Cove N. of Aberdeen A. 6. l. 14:8:56)  
*orthodesma Dickii*, <sup>(brown)</sup> ~~odontidea medialis~~  
*Fragilaria virescens*? *Himantidium*  $\approx$   
*Achnanthidium microcephalum*  $\approx$  <sup>(brown)</sup> *Diatomella*  
*varia* *waginensis*? and  $\approx$  <sup>(brown)</sup> *Diademella* (Lep.)

much build, lighter parts - best for sand & very dark  
*Fragil. virescens* ?

S.

412 (Aboyne, S side of river 15:8:56 Abt. 1)

*Naricula coccineiformis*, *Melosira* ? *systems*  
*Suriella linearis* (like *australis*) <sup>fragile, variegated?</sup> *adnata* *multifaria*  
*Eunoti* *tetraedra* (scapes), *Pinnularia* *Dactylus*, *nobilis*,  
*Gymnella affinis*, *Scotica*, *N. Vittula* *cervulus*,

413 (Steep behind Inn, Castleton of Braem B. 1. b. 16:8:56)

*orthotrichum* *orichalceum*, *adnatum*  
*australis* (scarce) *Fragilaria virens*? *Pinnularia*  
*interrupta* B. and a long variety; *P. nobilis* (very scarce)

for crude Lm 45g

414 (Pond behind Inn at Castleton of Braem B. 3. b. 16:8:56)

*Fragilaria capucina*, very abundant *confusa*  
 (frag. metolifera Rab.)

for crude Lm 260

S.

415 (Roadside on hill behind Castleton of Braem B. 4 &amp; 5: 16:8:56)

*Gymnella Scotica* <sup>Helvetica</sup> *affinis*, *explicata*, <sup>var.</sup> *Cleistothecium*  
*Suriella* <sup>var.</sup> *spiculata*, <sup>com. Helvetica</sup> *coccinea* *var. 1* Epithemis  
*infestans*, *ribbed*, *ocellata* & *viridis* (white), *Dickella*  
*Baltuerian* (scarce). *H. mankiana*, *odontidea* *borealis*,  
 most of *Grenvilleana* (Scarce), *Gomphonema vibrio*?  
 (very difficult to mount & requires long soaking in sulphuric)

416 (Roadside from castle of Braem up Glen Callader B. 7 16:8:56)

*Pinnularia* <sup>major</sup> *nobilis*, many species, some of the large tub.  
*Suriella nobilis*, *Stauroneis* *theanicornis*, *Naricula* *formosa*,  
*Gymnella* <sup>cuspidata</sup> *lappini*, *orthotrichum* *orichalceum* (*not* *Nernea* *distanti*)

417 (Surface of peat, N. of stream from Loch Ceander, Glen  
Callader. K. 144. 18:8:56)

*Suriella* *brevata*, *linearis*, *Pinnularia*  
*major*, *alpina*, *nobilis*, *viridis*, *semiflora*, *hirta*

Balsam very difficult to cut or - requires  
 a sharp knife

5.

- 418 (Bogs S. of the stream from Lock Ceandar, Glen Callander H.5 h)  
*Suirella linearis*, *Himantidium arcuatum*  
 ad *bidens* (very rare), ~~other~~ *erichores*?  
*Nardus stricta* & *Scirpus major*, ad *graham*  
 (very) *Pinnularia alpina* & *Luzula* (both very) bore.  
 - also (very rare) *Eunotia triodon* (very rare)

- 419 (Lighter parts of the last - 10.8.56)  
*Himantidium arcuatum* (small), *gracilis* (scarce)  
*Pinnularia borealis* (very rare), *Tabularia flocculosa*,  
 (good) *fenestrata*

- 420 (Bogs S. of stream from Lock Ceandar H.7 18.8.56)  
*Suirella linearis*, *Himantidium major*, *arcuatum*  
*bidens* (rare), *gracilis* (scarce), *borealis* (very rare), *graham*,  
*Cymbella* (not common), *luteola* (very rare), *Cymbella*  
*section*, *epitaxis*; *N. trichia* common  
*Navicula* *crassincisa*

redundant in later on S. 418, but with  
 more of *Him. bidens*, and *Cymbella*

5.

- Hill  
 421 (N. of Stream from Lock Ceandar, El. 3000 ft. H.6 h. 18.8.56)  
*Pinnularia lutea*! *Eunotia triodon*!  
*Melosira nivalis*, *navicula rhomboides*  
 (many *Gyrog. ericetorum* in a small streamlet)

- 422 (same place as last, light parts - H.6 h. 18.8.56)  
*Melosira nivalis*! *Eunotia triodon*,  
*Navicula rhomboides*!

- 423 (Stream from Lock Ceandar, Glen Callander 18.8.56 - H.8)  
*Tabularia fenestrata* & *flocculosa*  
*Eunotia* - *Diadomea* (very rare), *Calothecia* (rare),  
*erecta*? *Navicula* *crassincisa* (good)  
*Coccone Thwaitesii*, *Melosira* *nivalis* (common)  
 and *distans*?

5.

424 (Stream from Loch Ceandav, Glenballad, N.g. - 18.8.56)  
 nearly same as the last - but with more Eunstic.  
 tetrads. among heavy parts - also Hamat. areas.

5.

427/Loch na Gar, heaviest density G.s.h. - 8.56)  
 Hamatidium oxygen

*Hericium erinaceum* S.30.9. - in C.F.  
 Leng Flora no. 5.423  
 423 (Stream from Loch Ceandav, N.g. 10 - 18.8.56)  
*Tabellaria fenestrata* & flocculosa  
crude - same place as 8423.

426 (Spring w. Coreimulge 3 m. W. of Castl. Braem C.I. 19.8.56)  
*Trajularia*. *viridescens*, and *capucina*? / an  
 A. var 7. viridis odintz. meadow, Tabell.  
*lana* *fenestrata*.

428 (Loch na Gar, medium density. G.s.m. 8.56)  
 Eunstic. tetrads! Hamat. oxygen, (Hericium  
~~coccineiforme~~ <sup>in 8.676</sup> *Melotisca nivalis*, naevula  
~~longipinnata~~ <sup>in 8.676</sup> *coccineiformis*, *Gomphoneon*

429 (Loch na Gar, lighter density G.s.b 8.56)  
 Naevula *coccineiformis*, *Melotisca nivalis* & *dibam*?  
*Cymbella* *Setigera* & var., *Gomphoneon*  
*Pinnularia* *gracilis*?

S.

*Very fine material ready - and  
therefore S. 438 is lost 8m.*

- 430 Rock, n. of Cove near Aberdeen 1h: 8:56 A. 361)  
*Denticula oblonga, Cocconea Thwaitesian, Cym.  
 bell. ventricosa, Cocconea cistula, cornuta  
 Achmantidium lanceolatum, Odontidea mela-  
 lile tree (see ~~S. 4481~~ S. 4481)*

Same rocks as S. 4481, but at different place

- 431 (Teignmouth, side of millrace. Rev. N. Cope) *Navicula synchrocephala &*

- 432 (Glen Fincastle <sup>near by Lock</sup>, Killicrankie - R.K.G. 246)  
*Cymbella Ehrenbergii, cuspidata (var) affinis  
 helvetica, Lutica and var., Staurocione punctata,  
 Navicula firma var.*

S.

- 433/Bog near Lock Turnnel - R.K.G. (110)  
*Melosira nivalis, Cymbella ciliata? Lutica  
 affinis, helvetica, <sup>gracilis,</sup> <sup>runshi,</sup> <sup>fibulae</sup> <sup>and helvetica?</sup> Leptothrix <sup>very thin</sup>,  
 Gomphonema dictyonum, Himationium curvum  
 Cidens (var), Racoti (very rare), Epithemia <sup>suspecta</sup>,  
 Pinnularia <sup>large</sup> sp., Limnella, Ctenella, diatoms, etc.,  
 acanthella sp. interrupta, & -*
- 434 (Wattnean deposit, Caithness M. Peacock)  
*Heavier parts of S. 328 - This contain  
 a good deal of Nav. obtusa! if just on thickly  
 also Cyathella opercularis. The Epithemias  
 are less*

N.B. <sup>Mostly</sup> Some of another gathering at same place called S. 434.

- 435 (Wattnean Deposit, Caithness M. Peacock)  
*Navicula obtusa (Cope's name)? No. sp. not found  
 Epithemia ovalata (var.) Cyathella opercularis  
 antiqua (<sup>very</sup> ~~large~~ size), Oltotius arenaceus*

*Heavy parts of S. 438, but very much elutriated  
 for sake of the Nav. obtusa, which may be got 100 in. 100*

5.

- 438 (Hull, near ~~Leeds~~<sup>Per  
N. Yorks</sup> & ~~Leeds~~<sup>South</sup> Hotel) S. N. 208  
*Hormidium sigmoides*, *Nitzschia angulata*,  
*Pinnularia Johnsonii*, *Norscia Smilodon*  
*Actinaria*? (minute form) = *N. elliptica* Sm. mth) Nav. le-  
- infusoria var?  $\oplus$

- 437 (Minehead, Somerset - Miss. I. gifted her half 2 stages)  
*Podosira Montagnei*, *amphibiae antitelluriana*,  
*Cocconeis scutellum*, *Abbondanum adriaticum*, *minula*  
 (not my scars)

- 438 (Waithean Marl deposit Caithness - Mr. Peacock  
 Lighter parts of 435.  
*Mastogloia brevilli* (copious), *Spirula* - *Cyathella*  
*reticulata*, *Cyclotella opercularis*  
 (Epithemia nearly absent)

N.B. There were two samples of the Waithean deposit.

S. 328, there was little of *Cyclotella opercularis*, which was gathered  
 thickly. Since S. 328 and part of S. 434 was from the one sample,  
 S. 434 (rest), S. 435 (my reply) and S. 438 (light) are from the other

5.

- 439 (Site of Peas at Neyland L. Okeden. OK. 322  
*Nitzschia lanceolata* ??

very much smaller than Smilodon and more like  
*N. circula*

- 440 (Hull ~~near~~<sup>Per  
N. Yorks</sup> & ~~Leeds~~<sup>South</sup> Hotel) OK. 311  
*Nitzschia lanceolata*? (exhausted)

- 440 (Near Ballygawley N. I. May 1860 - S.N. 235)  
*Lymnaea adriana*, but thine too distant (about 20m off)  
 and does not enough infusoria - no Haworth in middle.  
 also *S. oocyathophora*, *Melos. varians*? In

- 441 (Pilling, Lancashire - C.J.  
*amphora hyalina*! *Pinnularia Johnsonii*!, *Am-*  
*phippsia* <sup>ataca</sup> *paludosa*!, *Spirula gemma*, *Nitzsch*  
*ligna*. *Lithiona* but more slender and  
 more distinct, steeper) *Tublionella punctata*, apically

crude  
 The same gathering yielded the rest!

5.

- 442 (Pilling, Lancashire C. S.)  
*Amphora hyalina* (scarce) seen on Bilean Amphora  
*alata*? *Pinnularia dolomiti*? (scarce) *Spirorbis gemmifera*?  
*Nitzschia ligula*? *Tyblionella peniculata*? mes.  
*griseata*, apiculate, acuminate  
 By breaking the amphora has become sp. alata.

- 443 (Haverfordwest, S. Wales, T. Okeden OK. 327)  
*Homeocladia fitiformis*? L. 8 mm. part

for crude see 464

- 444 (Haverfordwest T. Okeden OK. 328)  
*Achnanthidium lineare*, *Collotoma*  
*radicans*, *N. T. Z. lineare* var.  
 in OKOR lith 77 fig 1867 L. 5.60

not typical & to be tested in the ~~and~~ gathering  
 for the Collotonem in heavy parts in  
 L. 460 - See also 494.

S.

trace effects and *Henningsia*  
*D. tenue* var.

n. 251

- 445 (Hermits Well Redesdale OKG. 8: 56)  
*Amphora* (good) *spiculae* (small), *Cymbella* *apicula*  
*obtusa* (small), *Cymbella* *apicula*  
*obtusa*? *Odontidea* *Harritonii*? *multistriata*? (§ 1)  
 (What is called regular here must be on the C section,  
 few with thin ends curved - and large ones, angular  
 narrow ~~thin~~ ~~thin~~, and broader, ~~wide~~ at the ends)

and 9/11 AM

- 446 (Corrighill about 1 mi. S Corallieffie OKG. 242)  
*Amphora* *lepidopteroides* *Ralfsii*? *Amphora* *lepidopteroides*  
*Ralfsii* (good) *Spiculae* *stauroneis* *pulchella*, *Spirorbis*  
*buculum* (good) *Spiculae* *varia* (good)  
*Spiculae* - *Planoisigma rigidum* (very scarce)  
*Spiculae* *planum* *Ralfsii*, *Spirorbis*. *Staur. pulchella* (good).  
 446<sup>3</sup> *Spiculae* *T. P. longa* with some *Calymene* *distorsa* - others very good  
 447 (Glen Farg, on rocks 15: 8: 54)  
*Cymbella ventricosa* var ?? (in pub. L. ma-  
 culeata ?)

448 (Glen Long)

*odontidium mutabile* ? (small)

- M. Rylands *Merlus* this is in my book  
 (19 Oct. 1861) and that it is also in S. 430,  
 245, and 318

449 (Forth and Clyde Canal, A. Kennedy  
<sup>ventriculus</sup>  
*Cocconema gibbum* Thoms. = (*C. cistula*  
 var.?)

450 (Teignmouth - Stomach <sup>"allus."</sup> *gasteris*. Ad R. Caffell)  
<sup>(in fragments)</sup>  
*Eucanthe-godraccus* Diddalphi; Bailey  
*Tricardia altissima* all very scarce.  
 also *Syphlonella contracta*, *Orthodium maximum*

N.B. The *Eucanthe* require the autumn day.  
 some of the others in fall, at the slightest heat.  
Crude - published in S. 465

5.

451 (Teignmouth Rev. R. Caffell)

*Synodus Gailloni* & adsp. *Grammatophorus*  
 (thin & fine, ann. S. 639 & 3.307)

452 (Ecomouth Miss Cutts for Miss E. Hodges).  
<sup>Scutellum var.</sup>  
~~Brachionus~~ <sup>Scutellum var.</sup> *Lymnaea Gailloni*, *Grau-*  
*tellus marinus*

453 (Cerithus <sup>or</sup> Dunnet, recent G. P. Peach)  
~~Brachionus~~ <sup>Scutellum var.</sup> *Epithemis*.  
*gibba*, or *amphipleur Gibbonsi*,  
*Morularia gibberula* Maytaglom Smith &

5.

454 (Castricum, Brakkegroedepark. C.W. Peet.)

See S. 493 & S. 494

455 (Hull - Z. Okeden OK. 332 )  
*amphora tenera* ?! *Mitzaezi*. zigm.  
*Melvita nummuloides* Pn - See S. 488  
 Art. 20 found on m. 582

dead up - d remontant

crude

456 (Hull - Z. Okeden (B.R. 14) OK. 334)  
*Coscinodiscus sticticus* Ph.?

got by trawling some mud at horizon to water landing  
 from a Rotterdam vessel.

5.

ditch at Skirnlaug near G.N. Sper  
 457 (Hull, Fresh water f. Z. Okeden - OK. 344)  
*Mitzaezi dubius*? (*sticticus distinctus*)  
 (Same as in S. 103, S. 388, 296 and 520-2)

crude

458 (Haverfordwest Z. Okeden OK. 324)  
*Cocconeema flaccidum*.

for crude see 465

459 (Neyland, Z. Okeden - OK. 320)  
*Pinnularia cyprinus*, *Stauroneis salina*,  
*amphipora lyngbya* vibrea

460 (Haverfordwest Z. Okeden OK. 328<sup>b</sup>)  
*Pinnularia radicans*  
*Bellitorina neglectum*, *Spongiforma*  
*capitatum*? *Nitzschia*, *Lenticularis* & c.  
 From Okeden's bird S. 444

Many parts of S. 444 - same place as S. 491  
 but the <sup>is all</sup> ~~here~~ here <sup>there</sup> is  
~~here~~ <sup>is</sup> ~~there~~ <sup>here</sup> ~~there~~ than the S. 42. same

461 (Haverfordwest Z. Okeden OK. 329)  
*Nitzschia Pales*

seen off from 243 -  
 In open air N. J. lake. In. fm. Polus. L. 1. 1.  
 in fine condition. A few of them <sup>several</sup> <sup>large</sup>  
 form of *Nitzschia* but they are <sup>not</sup> ~~not~~ <sup>not</sup> ~~not~~  
crude

462 (Neyland. Z. Okeden OK. 337)  
*Cymbella ventricosa*? (small flaccid in S. 340)  
*Spongiforma tonelli*

The Cymbella seem the same as in  
 (OK. 276 =) S. 340

463 (Haverfordwest Z. Okeden - OK. 272)  
*Spongiforma* <sup>explicata</sup> ~~lancea~~ (plushful) - <sup>spores</sup> -  
*Lenticularia* <sup>affinis</sup> with a few of *S. lancea* -  
 also *S. radians*

464 (Neyland. Z. Okeden OK. 345)  
*Nitzschia lirostrata*, *lanceolata*?  
 (Same as in S. 439)

465 (Seignon by Rev R. Grewell)  
*Biddulphia* Bailey <sup>(Scarey)</sup> + *orthotrichum*  
<sup>coccinea</sup> <sup>dictyon</sup> (very small)  
*Trieurostria alternans*, *Coscinodiscus* <sup>centratus</sup>  
ovalisides? fm. - *Diatome* <sup>veratrum</sup> R. L. A.  
 \* B. Bailey letter in S. 470

From the stomach of *Phallusia* <sup>mustica</sup>  
assimilis - slightly  
 dried - due for balsam. well soaked for the  
*orthotrichum* - see S. 450

5.

466 (Penzance) J. Ralfr nov. 1856  
*Ecteinascidus subtenuis* Ralfr in Gregor,  
 dist. of Clyde - Synechia Baculus W.G. (= *S. pulchra*  
 var).      . . . . .  
 "        crude

467 (Dartmoor) rev'd by Gmelin  
*Melosira distans*?

Dust of sand with many bristles in Carb. of Br.

468 (Cornwall, arran - Beary)  
*Euphanes*, Ralfr! *Sauvagesiella pulchella*  
*Naria* var. <sup>didyma</sup> *coralloidea* <sup>marina</sup> *marina*.  
*Cocconia* *disrupta* W.G. - amphipr. *Lipidoph.*  
<sup>1</sup> *N. didyma* is much more finely studded than *Sauvagesiella*,  
 and this *N. coralloidea* more so from little *didyma*      S. 469  
 got by washing coralline specimens      13274  
 783.72

8

469 (Cornwall, arran - Lipster)  
*Sauvagesiella pulchella*, *Naria* var. *coralloidea*?  
~~*Amphor*~~ <sup>marina</sup> *Amphor* *Setosa* (= *A. effusa* var?)  
*Suriella festuca*, *Amphiprora* *Lipidoph.* *var. 5*.  
 Lipster parts of last (A. Lip. letter no 846)

470 (Cornwall) Lipster (part)  
*Coscinodiscus minima* Kutz from Sm!  
*Amphor* <sup>marina</sup> *Setosa* - *Canthigaster* <sup>rostrata</sup> *rostrata* &  
<sup>similans</sup> *parvula* (nugatana) - *amphiprora* *lutea* (Olney  
 Scarr)

471 (Silverstone) <sup>Heddon</sup> *Amphiprora* <sup>W. G. Huds</sup>  
*vitrea*.

472 (Neyland F. O. Klem. ph. 343)  
*Amphiporus nitens*  
 Buried up & remortised.

Buried, bark much broken up  
 from end in 467 (from end in 5.356)

473 (Cave at Cove, Aberdeen - ~~1950~~, D. Redfern)  
*Orthotrichum spinosum*, D. Redfern. *Achaea*  
*Thlaspi microcephalum*? *Ptyblionella canina*  
 Buried up. - regenerately on Cleft of Gable  
 near of the cave, at its further extremity. - new gamma 30  
 from end of 1950 spin. in 468

474 (Corriehill - from part of Corall. spic.)  
*Stauroneurus pulchellus* L.V. *Navicula didyma*  
*Amphiporus leptostoma* (very scarce)  
 None from last time there (32 or 35) - another violay  
 to water and under

almost same as 5.168

475 (Corriehill - from roots of Cr. off D. Gregor, g.d. 5)  
<sup>complex</sup>  
*Amphiporus* ~~leptostoma~~ (very rare, and easily broken), lepi.  
 doptera very scarce. *Stauroneurus pulchellus* common.  
*Eupodium pulchrum* (common)

476 (Corriehill - from roots of Cr. off D. Gregor, g.d. 10')  
<sup>area</sup>  
<sup>complex</sup>  
*Amphiporus* ~~leptostoma~~ (~~leptostoma~~). lepidoptera / very  
*Campulodiscus* common (& parrotia leaf)  
*Brachythecium acutum* (very scarce) may be *Brachythecium*  
<sup>(= hylocomium W.G.)</sup>  
*Luzula trichotricha* (= L. fuliginea)  
*Cocconeura* ~~discrepans~~ W.G. (= C. d. upha W.S.)

477 (Teignmouth - Stomach of *Phallusia sordida*)  
*Eupodium pulchrum* (some large), cr. of sun.  
<sup>bulbous (very rare)</sup>  
*Ralfnia*, *Coscinodiscus centralis*?, radiata,  
<sup>Bidens triplida (scarce)</sup>  
*concinna*, *escutifera*, *Populus maculata*  
*variegata* *Hedysarum* & *Lithospermum* (both *lithospermum* rare)  
<sup>& Amorph. arca W.G. 1/very rare</sup> *Urtica dioica* (scarce)  
 tom (magazine)  $\triangleright$  *Matthiola* (in 469)  
 di 2 Heaviest density - much bark

5

478 (Teignmouth - heavy density d. 5 )  
 sand. like loam or loam, but with ~~few~~  
 of Biddulph's Bailye - also pieces of broken  
 flint. Calcareous, iron pyrite (very rare) Nav.  
 Triangular plate here (May 1906)

479 (Teignmouth - lighter density - d. 10')  
 Coarse sand with scattered cl. (also broken  
 - alia, concretions, and ~~extremely~~ <sup>occasionally</sup> rounded?)  
 Triceratium variable, Brightwells, Biddulph  
 Bailye

480 (Teignmouth. lightest parts. of loam d. 20')  
 Detritus a. reparation s.v. // Corals, sand,  
 concretions: Triceratium variable, others

5.

481 (~~Limestone~~ - nov 5-6. Mif B. Hodges)  
*Sabellaria penetrata* on

481<sup>2</sup> *Columbella* N. Henratty

*Schizomenia mucronata* Sm?

The crude, heavy, very delicate one and amorphous  
 to some extent, used for fluid shells - Some burned in  
 cores are marked 5.481<sup>2c</sup>

482 (Premny Peak - 1848 Heavy C.T.)

*Orthidea orichalcea* (Smooth?) Tongue  
 cephalon, F. verrucosa. Liver. bilobata - Penn.  
 acuminate, nodose, evolute. Sm. (as to thickness)  
 evolute Sm (as to form = P. bilobata large) P. bilobata  
 P. evolute 125 darkish, notched, elongated, long neck?  
 slightly different from S. 120 & 5129. by hairy

*Fragilaria capucina*; also F. menziesii

483 (Same - light : c.t.)

S

- 484 (Pembrey Peat, Nov? 1854, Rev. J. Weston  
*Pinnularia alpina*, dactylus  
*Hornung major*  
*Euroth. tenuirostris*

nearly same as 144, but differently prepared

- 485 (australis (Wentworth <sup>has rock</sup> & K S 255)  
*Epithemia gibba*, ~~Zelkova~~, <sup>z.</sup> rugosa  
*Sciatomella Balfourianum* (var.)

- 486 (Mud in tidal ditch, Stavrosf. Devon dev'g Cuffell 1:57  
*Nariula cryptosticta*, <sup>amph. specie</sup> elegans, <sup>Westii</sup> pectinata, <sup>Westii</sup>  
*Nitzschia bilobata*, *Ligia*, *Tryblionella gracilis*,  
*Scutellaria marginata*, & *Pinnularia effusa*  
 intergrae (very rare) & *Pinnularia effusa*  
 var. *Juniperi* contracta

S.

- 487 (Tidal stream between Stavrosf. Dawlish Warren Devon)  
<sup>specimen var</sup> <sup>1:57</sup>  
*Fragilaria crescens* (Small), melior, flabell.

- 488 (Hall-) I.O. - ok. 358  
*amphora tenera* en S. 455

Mud, sand & silt 502  
 var. crassa

- 489 (Pichincha, near Quito, Peru - Dr. Gratiot n. 249)  
*Nariula* (same as in Long marine)  
*Nitzschia amphioxys*, ---  
 dried up & rehydrated

S.

490 (Baerlatten Brug  
Haverfordwest, OK. 361 2157)  
*Meridion constructum*

491 (Haverfordwest. OK 362<sup>a</sup>)  
*Homocladia piliformis?* *Gomphonema*  
*capitatum*, *Colletotrichum neglectum* - *Achna*  
*lineata?*

same as 494 *Ciliatostylus* and  
possibly place as S. 460 & S. 464, but the <sup>Homog. or</sup> <sup>diff.</sup> ~~nit.~~ seem,  
492 (Gare de L'Isle, Pyrenees - Prof Lorki)  
*Naiadella pyrenaica*, *Scita*, *complanaria* in  
form, *Succowia lacerata* <sup>cantabriensis</sup>

S.

Warton  
493 (Brakegroe deposit, Earlsfry  
*Naiadella obtusa*, *Peltula*, *etc.*  
*Cymbella Ehrenbergi* *a.*

The sand much, a different sample from S. 454  
coarser parts

494 (Warton  
Brakegroe deposit, Earlsfry - Light parts)  
much the same as last, but cleaner

495 (Wall of W. Norman Vicarage, Hull - ~~OK 367~~ 12) G.N.  
~~large~~ <sup>large</sup> *appendiculata* & *Pinnularia gibba*, etc.  
nature, <sup>red</sup> *Amphora Normana* <sup>Red</sup> *with* S.V.  
*capitata* <sup>greenish</sup> *earum* <sup>as to the valves</sup> *minutissima*

break up & grow the  
a new supply from Norm  
crude? requires the burned for balsam  
~~not for balsam~~

5

496 (Haverfordwest 3:57 OK. 387<sup>b</sup>)  
~~G. adesmiae~~ ~~|||~~ in sandy soil & silt  
 Himanthalia acuta (several fragments - S. 342)  
 Syringularia tenuis  
 pale soft algae by aid of ammonia  
 (then rehydrated?)

for crude see 495 496

497 (Hull (Dairy Creek) OK. 367 <sup>S.A. 25 Jan</sup> 102)  
 Amphibryozoan construct - ! Nitrochla. bilobata,  
 Liguaria sp. - author known <sup>in herbarium</sup>  
<sup>2</sup> <sup>referred by Dr. G. Norman to</sup> <sup>as</sup> <sup>and</sup> <sup>by</sup> <sup>3. 1. 25</sup> <sup>herbarium</sup>  
 497 <sup>the</sup> <sup>lithion</sup> <sup>only</sup> <sup>all</sup> <sup>used</sup>; the remains were mixed  
 with G. Norman 102 from Victoria. Drift Hull. What the  
 not cut N. t. bilobata - as a <sup>green</sup> <sup>greenish</sup> <sup>leathery</sup>  
 Engbl. <sup>Scutellaria</sup> crud - for crude see next

498 (Hull OK. 367)  
 Amphibryozoan construct - various forms of *L. c.* |||.  
 Nitrochla. bilobata, Liguaria sp.

5

499 (Dockyard at Pembroke - OK. 380)  
 Amphiplexa ligulata! Pinnularia. <sup>Stom</sup>  
 - cori? Achanthia longipes. Schizoneura concava?  
 Smithii. Nitrochla. angularia.

500 (Neyland OK. 379<sup>a</sup>)  
 Schizoneura helminthosoma

sent as "L. divergens"  
 for crude see 502

501 (Hawkingshaw Ph. of Graham S. E. of Slapton 28:3:57)  
 Saccorhiza cistula, and cylindiforme

5

far down right water mark

- 502 (Hanging shaw <sup>cathart</sup> of <sup>new ligament</sup> Gobek. S. 9 1889 Aug 28. 8. 37)  
*Nitz. tenue*! - *a dubia* - *Harrach amphistom*  
*Syphlonella granul* (extremely scarce)

In diagram of *N. tenue* in *Lyncea vermicularis* 115

- 503 (Hanging shaw <sup>new ligament</sup> nearly same place as last)  
*Nitzschia tenue* and *odontostoma* parasitic,  
*Syphlonella granul* very scarce

(lighter density of the rest)

- 504 (Hanging shaw same place as last)  
*Nitzschia tenue* with a few frustules of  
*Syphlonella granul*! (scarce)

Heavier density of last  
 for crude see No. 503.

5.

- 505 / Mouth of Teign. S. Devon. Red M. Crayfish  
<sup>N. granul</sup>?  
*Nasicula palpebralis*? minute & <sup>near</sup> <sup>large</sup> <sup>size</sup> <sup>and</sup> <sup>shape</sup> <sup>as</sup> <sup>in</sup> <sup>Cumbria</sup> 1866  
 able very <sup>scarce</sup>, *Nitzschia* <sup>very</sup> <sup>scarce</sup>? *Tecula*? <sup>Scutellaria</sup>  
*Melosira Westii*, *Pleurosigma transversa*

For crude (light) see 504

- 506 / Exmouth? with cult. for Miss Hodges  
~~Stipe diffusa~~ ~~Solmaticea~~ <sup>or</sup> ~~coarse~~ <sup>thin</sup> <sup>long</sup> <sup>rigid</sup>  
 valves tawny, twice distinct shape, - there is  
 a thick grey <sup>yellow</sup> <sup>also</sup> stipe, to my h. <sup>also</sup> Stipe. <sup>coarse</sup> <sup>thin</sup>  
 crude for burning on corners

see S. 529

- 507 / From a Mollusk shell <sup>on</sup> <sup>366</sup> <sup>edge</sup> <sup>of</sup> <sup>shell</sup> - *Hydro*  
<sup>Bedrock</sup> <sup>Bailey</sup> - Diatoms vegetation <sup>Red</sup>  
 (same as in S. 1. 50 + 3. 465. fm  
*Neurocolema lanceolatum* Norm. in Ann. n. H. XX. p. 159  
 (Norman 1879)  
 (Some *Nitz. tenue* had been in the later  
 when last me.)  
 crude - the rotted if for bottom

S

508 Neyland  
*Achnanthus pauculus* OK. 390<sup>3</sup>

only slightly boiled for 10 min. Blown later  
for crude see 506

509 Neyland  
*Achnanthus pauculus* OK. 390<sup>3</sup>

calcined - for crude see 506

510 (Hull

*Nitzschia*


all N. T. Z. dubia. Linné says. (names  
publ. was in the latter previously) Oliver specie?

510<sup>2</sup> (Ulverstone Mif S. Hodge)

*Cymbella Helvetica* (no stipe)

*Diatomea* <sup>terrea</sup> ~~terrea~~ (then D. inflata) (no stipe)  
*Leptothrix* <sup>terrea</sup> ~~terrea~~ (very small) (no stipe)

N.B. The Cymb. is very like common cymbiform



S.

511 (Cumbrae R.H. 4:57) "  
*Dickelia pinnata* (mixed with Schizone  
leibnitiana)

512 (Cumbrae, no care, R.H. 4:57)  
*Navicula tecumida* Judd

513 (Wall in Haverfordwest 4:57-OK 394)  
*Achnanthidium eravestatum*! orthos  
mirabilis! *Navicula pusilla*! *Nitzschia*  
amphibia!

all as in S-245, but rather more boiled  
and with more of carbon

dried up and remixed - eight  
hrs boiled in 500

5

514 (Haverfordwest. Z.O. 4:57-

*Synedra affinis?* (very minute)*Merdion circularis*

Forerunde der 507

515 (Neyland Z.O. 395<sup>a</sup>*Schizogonium implicatum*

billed for erode der 508)

516 (Rock at Neyland) F.O. 391<sup>b</sup>  
Gemmig ~~and~~ <sup>and</sup> ~~luffatg~~ var. <sup>2</sup> ~~omphalothecum~~ <sup>1</sup>  
~~Dark green~~ ~~dark~~, Nit. linearis.

for crude der 505

5

517 (Gourock. on Esteromph. April 57 A.H.)  
in rocky pools, sea shore  
*Nitzschia*

(an epiphyte?)

same as in 18

crude -

518 (Gourock on Esteromph. April 57 A.H.)

*Nitzschia*

same as last, but boiled for 1/2 min. in acid.

Invisible in balsam - only for dry morning

519 (Neyland F.O. - OK. 400 - June 1857)

*Synedra affinis*, Gaillons, Bidensphoe  
aurita, melosira nummuloides

S

520 (Neyland, a circular pebbly rock. OK. 399<sup>\*</sup>)  
 Schizoneura pustula?  
 Nitzschia Clotellum! sigma &

523 (Ex mouth of Cutler River 54)  
 Odosphenia Lyngbya var. s.v. very narrow  
 valves crystalline - stipe scarcely more than 36-38

crude (fertilized in nest)

521 (Neyland  
 Schizoneura  
 Nitzschia sigma  
 OK. 399<sup>\*</sup>)

524 (Hill behind Lambsk, area A.N. July 57)  
 Ooids or ichthy! Triglochidium avicinum  
 and - Pinnularia Tabellaria, majority brown

ended

522 (Spring in Strath Tummel - D. Greville "248")  
 Narcole crenulata? Pinnularia longi-  
 tenuis (var) fr.

This was much broken and, and still the rocks  
 under water only held 2 dredged w Turf. - In mud  
 on 512

525 (Campsie Glen - A.H. 3:8:57)  
 Cymbella scotica? (var) - Nitzschia tenuis  
 (straight, & small), Cocconeis v. sp., dentata  
 v. sp., Cocconeis Thwaitesii

\* S. m. conspicuum - is it only a var of N. rhomb.

S

- 526/ *Torrey* - on *Callith. Burrieri* on  
Grammatophor ~~near~~<sup>near</sup> ~~Leptodeira~~<sup>Leptodeira</sup> 110 m.  
investiens

thin form of *G. marina* less lanceolate than *G. septent.*

- 527 (*Torrey* - on *Call. Burrieri* I-T.N.)  
come on the above, coerulea

- 528 (Exmouth. mif Cutler - 8:57)  
Grammatophor subtiliſimis Bailey?  
red up - removest

thinner than *G. marina* - about  
the same as *G. macilenta*, which may be only -  
narrow variety.

S.

- 529 (Buddleia Salterton - May 1849. Mif Cutl.  
*Podosphaeria Purpurea*

values changing  
still more than in S. 523. but not faint.

some bits of the eggs mounted in fluid are S. 529\*

Stipe and stalk well - this is 8:57 - see S. 506

- 530 (Well at St Andrews M. McDonald. 8:57)

*Syndra hamata* var. radicans tr.  
purely fresh water

on *Vancheria*, a cladophor. - for crud. see 518

Form of *Turritella* as in *S. radicans* B. *Micromesistius* about  
36 or 38 m. (finer than in *S. affinis*)

- 531 (Montland Canal Glasgow 14:8:57)  
*Syndra ulna* - <sup>c. hamata var. radicans</sup> L. as in 530, also radicans  
var. - *Diatome vulgaris* tr.

on *Vancheria*, a cladophor  
for crud. see 516, 517

S.

532 (Hull -

OK. no 368)

"Cocconeum with broad turned up" = Cymbella  
 helvetica = *C. plicata* W. Gray?

533 (salt marsh at Havenfordwest over var. - OK 393)

Spirula amphioxys (fond) - Nitzeck  
 Alvea! - *Sertularia elongatula*, var. fusca

534 (Hull

Nitzeckia vivipara <sup>GN/see</sup>  
 granular, Spirula amphioxys.

See 5.721, #591

S.

535 (Sidmouth arcticum, Adriatic <sup>mp</sup> Buller /  
 Melobionema ~~adriaticum~~ <sup>mp</sup> Synder. Gaillouet.  
 maxima (Gardes - see in 5.79 (5.399)  
 Grammatophora Lepthostoma! Podoseta montagnier  
 (see in F.V.)

535\* a few slides were kept with much of adriatic!

536 (Sidmouth - from Gladstone Bay <sup>mp</sup> Buller)  
 Podoseta montagnier - below F.V. (?) - also  
 Ababdon arcticum, adriaticum as in 535

nearly crude - for burning on cover, & bellarmine

537 (Blapham, York <sup>(= baculum wsg)</sup> C.J. I)  
 Stararia Leite W.L. - Cocconeum cistula  
 Synder & adriaticum. —

This does burn the burned on the cover, and  
 the bellarmine - on fire is <sup>diluted</sup> held in the  
 water

S.

538 (Dunfermline, eyrethrin Mc Macdonald)  
*Epithemia turgida*, *Lorenz Gibb*, ventricosa

539 (near Ventnor, Isle of W. 1851) J. Norman  
 Grammatophyton <sup>earlier page? (three lines)</sup>  
~~see S. 307~~ <sup>see S. 307</sup> ~~see S. 307~~ ~~see S. 307~~  
~~see S. 307~~ ~~see S. 307~~ ~~see S. 307~~ ~~see S. 307~~ ~~see S. 307~~  
 serpentaria (stein) - Cichor  
 longipes, Synidre <sup>nitella</sup> <sup>nitella</sup> <sup>nitella</sup> <sup>nitella</sup>  
 but ~~the~~ north & three about 58 or 60 m. (00), Podosphaera  
 n. Thunberg (very rare)

540 (side of excavation at Woolwich, bonepit below  
 the Thames high-water mark; the water permeating  
 through the clay, at the entrance of a new dock,

- J. Norman - 29 Aug. 1857  
 Bicyclella <sup>nitella tubifex S.V.</sup> <sup>and Eudistoma small</sup> <sup>very rare</sup>  
 var. <sup>nitella</sup> <sup>var.</sup>

Nitella <sup>0/ minute - an N. less rare</sup>  
 amph. Salina L.

buds

F. Mr. T. G. N. Hunter Cyc. mimeticus var

5

541 (same as last -  
 as in last

J. T. Norman 29.1.17

finer parts, crude.

Some Podocystis from S. 21 has accidentally got mixed in this ad. 540

542 (Lewes M.S. 1853)  
 Bacillariae parasiticae

for living on corals & bivalves

543 (Brackish water, Lambeth - 22 July 57)  
 Navicula Rummeli, amphipora ~~leptophora~~ <sup>leptophora</sup>?  
~~leptophora~~ didyma - menard marum w. Z.V.

heavy parts - for light see next.  
 for corals see 523

S

544 (Brackish water, Lemurak - D. July 1857)

Nitzschia *meridion marina* (L.) M. B. C. H. (lithocidum)  
short leath / (an N. palm var?) - Fregil. thistle  
small (?) Cyclotella <sup>not</sup> (a. in S. 540), Navicula  
numerous, fragrant, sp. (a. in S. 540), Navicula  
light parts of leaf - for crude see 523

545 (among Marchanti. polym. in struc. of <sup>D. H. Sept 57</sup> Gouraud,  
*Himantidium gracile*)

546 (Brackish water, Boston, Lincolnshire - Dr.  
Moor - 9:57 "A. 2")

Pleurocione Spencer, Navicula cryptos  
capitata

crude

S

547 (among moss on trees in Pulney mountain)

Penins. of E. India.

Orthotrichum species?

548 (among moss, on top of well, near Partick  
*Pinnularia borealis*) <sup>19:9:57</sup>

549 (among Moss 39, fm trees in Peninsula of  
E. India (Dindigul?))

orthotrichum species?? (remarkable from  
with close awl) *Pinnularia borealis*, Nitzschia  
amphioxys, Navicula multistriata? (Staur. semin.  
shrub.) *Achnatherus ventricosus* (Duby) from

5

550 / <sup>letter, D. Moore</sup>  
~~Ascidia~~ <sup>as Herb Harvey sub nom</sup>  
 "Cymbella cyathiformis")  
*Epithemia hypnoides*, *Othonoia arenaria*  
*Cocconeum cyathiforme* small var.

551 (Gowrock <sup>10:57 A.M.</sup>) 10:57. A.M.)  
 - Cymbella Scotica var. angusta:  
*Naria* *trivittata*

552 (Gowrock - on Mop on a Plant Tree. Elevation <sup>200 feet</sup>)  
*Orthosiphon strobilifer*

5

553 (Penzance gull by washing shore sand  
 Oct. 1857. A. D. G.)  
 auf der Fron <sup>5. v</sup> <sup>s.v</sup> <sup>1</sup> *Pinnularia*  
*capricornis* var? (*capitata*)! Navicula retusa  
 ex rugulata sup.  
 for covers - (for balsander 5.50g)

554 (Teignmouth (Nettleton Point) Oct 1857. A. G. G.)  
*Actinularia peregrina* and var? nitida  
*bilobata*, *Achraea subcylindrica*

555 (on a Rock, among lichens, at Goatfield, N. L. P.,  
<sup>10:57 A.M.</sup> *Himantidium Soleirolii*? (in the normal  
 state - and without the internal septa)

5

556 (S. Andrews. East Bay - 3:10:57)  
*Schizogonium multifidum?*

for crude see 539

557 (S. Andrews, on Clad. superba 29:9:57)  
*Schizogonium laciniatum?* *Synedra* -  
 - - - (in nest)

for crude see 540

558 (S. Andrews, Bathing place 29:9:57)  
*Synedra horvathi* R. ~~thick~~ three very far  
 apart 2.5 mi. 00. Elevation unknown.

for crude see 541. This *Synedra* seems the  
 same? a. 66.

5

559 (S. Andrews Bathing place 29:9:57) -  
*Zonophorella marinum?*

Never confused w. w. *S. marinum*  
 found on an alga in Salt Water pools with rocks,  
 but *Dicotoma vulgaris* was mixed

560 (S. Andrews - Bathing place 29:9:57)  
*Synedra gailloni*, <sup>horvathi</sup> ~~superba~~ - ~~nest~~ (as  
 in 558)

561 (Tents near <sup>(4)</sup> S. Andrews, among Lyngbya ferrug. 1:10:57)  
*Pinnularia cyprina* (espirus) - *Navicula*  
*Westii*: *Achnanthus subfusilis*

lighter density of nest

562 (Tents meet (1) S. Andrew among Lycopodiums.)  
*Navicula Westii*, Linnæus  
*Pinnularia cyprinum* (Scutell.) - act. surp.

heavy density of lichen

563 (mop on Elm at Abercorn, W. Lothian. 10:57)  
*Orthotrichum nitratilis* (Sund.) - naviculae  
 mutica? gr

564 (mop on Elm at Millport, Cumbrae 22:10:57)  
 orthotrich. nitratilis, naviculae pusilla  
 mutica. *Pinnularia borealis* (Flotz). -  
 (little or no Kitz. amphioxys)

565 (washed from sand, near Millport, Cumbrae 22:10:  
 amphioxys. *Ralfsia* <sup>var.</sup> ~~lepidota~~ (Same as in 553) 2  
 met. ~~Microcoleus~~ Donk. ~~Microcoleus~~ <sup>met.</sup>  
*Phaeoceros ceterium* - Nav. *leptobrana*  
 lumenous. *Actinodiscus* <sup>griseus</sup> Kitz. *spissatus*  
 coronata. *nitidum* Grig. - *Navicularia rectangularis* M. S.  
*Actinodiscus* <sup>griseus</sup> Kitz. *spissatus* <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~  
*Actinodiscus* <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~  
*Actinodiscus* <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~ <sup>griseus</sup> ~~Kitz.~~

566 (Sidmouth on *Gelid. cornuta* trifidae  
*Grammatophorus tropica* <sup>microcystis</sup> (Conrad, a. 579)!  
*Podostemum* Ehrenbergi secur

567 (Caen - Prof. Chauvin)  
*Frustulia acerps* " <sup>and</sup> *itzschii*.  
 sometimes marked S. 567 <sup>longitudinalis</sup> <sup>longitudinalis</sup>  
 in the corner.

*F. acerps* of Huds. is united by him by Syndr. Garrel. but that seems composed of two species - one a "Syndr." - the other a "itzschii".

5

568 (Peru - Freshwater l. Ralpy)  
Homoeocladi. Sigmide

S.

571 (Teignmouth, on wood of Harbor, )  
amphiblenni. niflet <sup>New. O. Cupell</sup>

569 (Berlin deposit - Kutzay - see Shub. mix  
Orthocid. punctata <sup>tab. 14</sup> Chub. m / = Gallinella granularis  
lita. Mexicana? decaps. shrub.) - an.

## Cnidar

572 (New Zealand, from rocks of Janie  
Grammatophor <sup>undulata</sup> angulata, <sup>angulata</sup>  
Cnidariae <sup>undulata</sup>, hamatiform, <sup>angulata</sup>  
Cnidariae <sup>undulata</sup> ~~Cnidariae~~ <sup>undulata</sup> ~~Cnidariae~~ <sup>undulata</sup>  
Epizyguria ocellata W. C. + pulvilli  
Abdomen <sup>(campanula)</sup> (campanula) <sup>(campanula)</sup> <sup>(campanula)</sup>  
Bullia <sup>(campanula)</sup> <sup>(campanula)</sup> <sup>(campanula)</sup>

572<sup>2</sup> Euplectrichthys - from Polyozoa Harveya (adult)  
572<sup>2</sup> " " crude - (juvenile)

570 (Berne D. R. J. Schmidt)  
Diatoma ~~guttatum~~ Ehrenbergii

573 (New Zealand, from Bullia callinich)   
Euplecteria <sup>undulata</sup> ocellata W. C. - Gram.  
- mat. flor. angularis Ehr

for crude see 572<sup>2</sup>  
mixed in a small thing  $\Rightarrow$  not unlike.  
little Neomantidium, but with nodules - an

S

574 (New Zealand - n° 3879 Dr. Harvey).  
*Synedra Gailloni* B. Kutz

575 (Van D. Land n° 1893, George Town) Dr. Harvey  
*Grassoa et flora macilenta Sm?*

576 (New Zealand) Dr. Harvey  
 576 (New Zealand - F. Water? 2194)  
*Melosira subflexilis* Kutz? (vice Smith)

This has the char. of *M. varia*, but the opercular  
- and appearan of *M. subflexilis*.

\* calciated dry: for cover (chitinous)  
 \* a beltam

S

577 (New Zealand - F. Water? 2194)  
*Melosira*

Corolla

578 (New Zealand - F. Water 3832)  
*Synedra ulna* var - *Cocconeis circula* var.  
*achmauthidium lanceolatum* (Smell?)

579 (Murray Bay, N. Shore of Lawrence Mif. Bay 6)  
*Synedra affinis*. - *Podosphaera* *Ingenii*?  
 (there very delicate). *Melosira* remains  
 (as the other two flanges of life) and sporangial  
state (*M. dubia* Kutz). Denticle?  
 (minute).

very slightly build - requires the removal  
a cover or slide, before being observed

S

580 (Cumbræ - dredged - 10:57 - heavy)  
*Pleurisigma stigosum?*

581 (Cumbræ - dredged - 11:57 - light)  
 cumbræ - Navicula? ~~—~~ - very occurs  
*Pleur. stigosum* and an anomalous ~~stig.~~  
 - also, benthic grass, amphipods, <sup>Ralfson</sup> (S. 565)

582 (mop of Elm, near Dumfries 11:57)  
 Pinnulari. border chiefly of ~~the~~ <sup>old</sup> Hodder  
 species, miserable, Navicula ~~dentata~~? - <sup>care</sup> (S. 565)

S.

583 (Neyland 01.413 )  
*Achnanthes longipes* var. *valvis acutis-*  
*natis* (A. Brébier Kutz) *Grammatophora maci-*  
*lenta* and *subtilifera* (only var?). *Synedra*.  
 (as in S. 558, and 66). Schiz. Grevillei frust.  
 scarcely bristled; requires to be calcined - heavy.

584 (Neyland 01. 1013 413<sup>b</sup>)  
*Grammatophora macilenta* and *subtilifera*  
 - *Achnanthes longipes* var. *acute* - *Synedra*  
 harvey. (as in S. 558) - Schiz. Grevillei frust.  
 much bristled - heavy - fit for balsam

585 (Neyland 01. 413<sup>c</sup>)  
*Synedra harveyi* (as in S. 558) - *Grammatophora*  
*macilenta* and *subtilifera*. *Achnanthes longipes*  
 var *acute*!

light parts - ready for balsam

5

586 (Haldon, near Teignmouth - A.C. 11.37)

*Pinnularia hemisphaerica*

587 (Back Nettleton, Lancashire 10dm 1852 G.W.S.M.)

*Campyloodium sp. alr. (abundant)*

{ often green

In the fresh air very thinly, on it is very dirty

588 (From moss off a tree near station for S. 245)

*Rimularia bridgemanii* (OK. A. 14) *Navicula pulchra**Schizothrix excentrica*

{ greenish brown

S.

589 (Penzance

OK. 426

same as S. 553, but dirtier, some more  
brilliant and, and for balsam - S. U. of  
Navicula retusa are sometimes got

590 (Minehead, Somerset Mif. J. Gifford

*Nitzschia tenuissima*, *Plaeosigmo* *Sporenii*

+

same as 555, but brilliant ~~and~~

591 (Minehead, Somerset Mif. J. Gifford)

same as last, but coerulea, andwith more of *Tephromella Scutellum*??

(After T. gracilis var?) - Mrs. Balfour may

seen

592 (Sand on Silverton Mfthg) 5  
Amphorhydium

595 (New Zealand on T.W. edge)  
Epithemia v. sp.

see 5. 632

593 (Kotzebue Sound D. Harvey)  
Synedra affinis Podocia Hornemann large  
<sup>Ceratostoma & Thalassia</sup>  
= Cyclocaea subtilis (B.) and small - ~~Ababdonia~~  
~~crozieri~~ Shaff.  
arenaria Dr.

596 (Chinc (Peru) Guern m. Hemmig)  
<sup>(Antennulae & Oral?)</sup>  
Eupodium w. 2 holes Spatangoid  
arachne

594 (Petrovskoostrovski Bay, Katalashka D. Harvey)  
(same as last but without Podocia)  
Mabb. arcuata var (A. crenulata)

Harvest  
for Dr. D. Harvey Jan 2 1832 1861  
597 (Chinc (Peru) Guern m. Hemmig  
Spatangoid v. f.

L. Blatt

S.

598 (Washings of algae, T. &amp; meadow N. Zealand)

*Hyalodiscus* .... Shr. *Coscinodiscus* ....  
*climacophora* <sup>Petersen and</sup> *austrolii*. *Aulacodiscus*, 2 sp., (var.)  
*Triceratium* (Aug. var.), *Euphemocellatum*, <sup>and *fulvum*?</sup> *Dick-*  
*dulphi*.

598\*. a few slides of a second washing from other algae.

599 (Californian Green) J. T. Nor

*Aulacodiscus* <sup>merfordi? Dalf. & erosa?</sup> *coscinoides* *Dickdalia* *Juncea*  
*Lepid. scutpus* (*caerulea* Paul.). *Triceratium* *sarcinum*  
*Dicella* *variegata* (Lob.)  
*Biddulphi*, *Actinopeltis* *plumosa*.

Some slides marked 599\* are from an inferior preparation  
 in slide no 609

600 (Dawlish Warren Rev. R. Crispell)

*Coccinodiscus* <sup>var. in Rose</sup> *coscinoides* *levis* *Lev.* <sup>reducta</sup> *coscinoides* <sup>Lev.</sup> *from Daffin*

*Nanula* *pectinata* (*Stimpson*). *Smithia*, *humerosa*.

*Lites* (= *N. villosa* Daffin), *retusa*. *Stauronema*

*Zetellia* <sup>7.V.</sup> (= *N. Zetellia* Daffin), <sup>and *lutea*,</sup> *coniformis*.

*Denticula* *stauronema*, name.

\* In the Museum in (b) not defined as a new genus

S

601 (South African Green - C. F.)

<sup>Aulacodiscus</sup> *Petersii*, *Canthistera ornata*,  
*Triceratium* *comptum*, *grammatophyllum* *undulatum*,  
 601\* same but not so good

602 (Calvados 1852 M. Detribes) [

*Epithemia* *constructa* Daffin (non Sm.)

*E. Westermannii* var. *Sm.* (non Kutz.) *lutea*, *gibba*

[

603 (Cherbourg 1859 - M. d'Orbigny)

"*Coscinodiscus* *coccinoides*?" = *Epithemia*  
*subtilis* Gray, but striae very distinct in slide  
 after boiling in acid! also *Compsyph. Hedder*

S.

604 (Mexican Guano. D'Yandres)   
 Infusoria (○) *Heteroplaesia avastri*. Tricost  
 variable spm

This contained a large percentage of *Calyptothrix*, sand  
 & sand seems to have been an adulterant. The article con-  
 tained of this, and a very little *Pennaria* Guano

605 (Pilling Rev'd Mr Barnister per C.J.)   
 Pleurocystis Fasciata, angular, quadrat  
 lattices. *Hippocampus* (male var.) Survival  
 genus &  
 all easily solv'd in H<sub>2</sub>O dry.

606 (Deposit from Ardèche - Dr Breysse 25)  
 "Mollis ad ciliat. bordum" Kutz. = *Gallionella*  
 marcescens shr. = *Orthotrichum* marcescens

607 (Fabache - M. de Bréjison)  
 amph. exiles

608 (Fabache Chabaud - M. de Bréjison)  
 Surirella crumenaria

609<sup>1</sup> Patos or California Guano. (Combe Bay)  
<sup>not dried this mixed with 609</sup>

609<sup>2</sup> <sup>analogous</sup> <sup>inland</sup> with 5-6 mgs. of sand  
 hair spn - *Bidwellia*. *Luzula* (Bacterwash  
 <sup>kind of</sup> <sup>perox.</sup> *Hemidiodia* well  
 *fruticosa* & *Cryptotaenia*, among the lighter parts  
 Ocularis scutellae and *Penitularia*?

+ few fort. amphibiae and fossils  
 L. *Bidwellia* as above

S

610 (Great Meols, Cheshire T. Comberby)  
 Navicula (same as in 1332139)  
 also Nav. small. ♂ resemble *Hausmania*  
*Nitzschia virgata* R.

611 (Adamstown Bay, Jamaica, Washingtonophyte N.Y.  
*Cocconeis punctifera* Grun. in amphipod  
 ornata, *Pseudocystis americana*

Some T.W. Epithemia re-washed for the table by Dr. Grunow

612 (<sup>more</sup> ~~Adamstown~~ Bay - Cornwall - G. Norman  
*Eupodium pulvinum*, Syph. ~~concreta~~.

From the stomach of *Pecten maximus*

They were said to contain *Spatangus* & *flabellum*  
 but none ~~had~~ can been seen by me in this

S.

613 (Swansea, waterfall, valley beneath OK. (132) 431)  
<sup>(waterfalls in S. occid.)</sup>  
*Epithemia argus* Navicula ~~tabida~~ (few)  
*Micantidium bidentatum*, *Denticula* <sup>lancea</sup> ~~oblonga~~

614 (Swansea light parts of S. 613 OK. <sup>431</sup> ~~some~~)  
 Navicula <sup>tabida</sup> <sup>nigra</sup> ~~lancea~~, *Denticula oblonga*, *Cocconeis*  
*Thwaitesii*

615 (Swansea near the head OK. ~~7/11~~ 432  
<sup>argus, adusta, tenuissima, irregularis</sup>  
 Navicula <sup>argus</sup> ~~tabida~~, *Epithemia* <sup>adusta</sup> ~~argentea~~, gibba <sup>var.</sup> ~~tabida~~

+ spectra of *Prister* body, as the 2 sp. char.  
 it not of *Prister* which has no porinae, and  
 is unlike from *P. Zebra* - see 8.621  
 spectra of *Prister* body

5.

616 / Swanson  
*Synedra tenera* deformis, lunaris and distorted  
 var. ff

off 429 1

Mycorrhizal

st. sterile branched

617 / Godwinell

5/1/12

Forest flora 1000 m

620 / Heyland

5/1/12

Forest flora 1000 m

www.wwwwww

618 / Glance Rock

5/1/12

Forest flora 1000 m

621 / Heyland

5/1/12

Forest flora 1000 m

For the ~~other~~ diversity see 687.

600' - 650' <sup>and maculate</sup>  
annular or tubercles, small, <sup>and</sup> <sup>white</sup>  
white (scarce)

on same as S. 009, but smaller  
thinly <sup>irregularly</sup> <sup>white</sup>

600' - 650' <sup>and maculate</sup>  
annular or tubercles, small, <sup>and</sup> <sup>white</sup>  
white (scarce)

thin white

Swansea  
Epithemiidae, <sup>affinis</sup> <sup>18. 17</sup> <sup>Smith's Shells</sup> (not of his figures)  
Cyclotella Kutzingiana

very rare

1 Swansoo Bog  
Syndra pulchella, odontidea Tabellaria.  
O. nobilis (Smell)

very rarely

600' - 650' <sup>and</sup> <sup>white</sup>, ash. 11. 7.  
Leptothrix longula, thin white

600' (Swansea Bog)  
Leptothrix longula

rare <sup>scarce</sup> <sup>scarce</sup>  
scarce well started in her path.

5.

- 628 / London-hurst Dr. Waller )  
 ✓ *Hydrostoma compressum* not from S. V. of St.  
 Trigem - Achmatov's wood, Novocastrian  
 Australia?

5

- 631, New Zealand - off Balli. collitrich. C.S. 1  
 ✓ *Graconia caelata* n.sp. ~~E. f.~~, *Euplexia collata*,  
*Grammostola* v.s. *amphibia* [Yonaguni]  
 — ~~(Mother Gullies - see S. 573)~~

- 629 / Swanson Dock. Okan OK 63  
 amphipods, amphipods (ungsav)  
 Melvin numerous soft shells

Crust

- 632 (New Zealand - L Water? ~~near Hobart. T.~~)  
 : *Epithemia* v.sp. v.s.  
 nearly the same as S. 595, but more mixed

- 630 (New Zealand, off Balli. collitrich. C.S.)  
 ✓ *Aleurodiscus* (near formosus) <sup>new sp. S. 637</sup> very rare  
 Bridgeman . *Euplexia pulchella*  
 and var? (~~new~~, not ~~new~~) or n.sp., *Coscinodiscus*  
 ---

Slender stalk uniform as they were prepared from various  
 trunks, the residue of which from 6.630

- 633 (New Zealand) C.S.  
 orthotricha cricidalea, *Morularia pectinata*?  
<sup>earlier Wright</sup>  
 Curvata? " <sup>misnomer, probably</sup> amphipoda  
 On a shrub like *Ligustrum*, where the epiphytes mainly  
 sought the burned on cover & became  
 a little of S. 628 get mixed

S

634 (Swansea-Dock 2 October OK 165<sup>a</sup>)  
*Pinnularia Johnson Schizomerus* (large  
 and siliceous) -  $\text{OK}^2$   $n \pm 10^{\circ}$  ~~1~~ *Acanthopora*  
*contracta* var.)

635 (Utracomb  
Cornwall  $n \pm 10^{\circ}$ )  
*Tetracyclus lacustris*, *Aimantidium pectinatum*  
*odontidium mesodon*, *Tabularia flocculosa* and  
*fractata* sp.  
crude

636 (Vidzome Dark Mill G.N.  $25 + \text{OK} \cdot 367$  (as outlined))  
 amph. crustacean, Nit. 2. 18mm (libt. 8cm)  
 (tbl. granular 2cm.)  
 OK 367 was from Denny Carts, down ~~the~~ <sup>S.</sup> ~~the~~ 10<sup>2</sup>  
 but was accidentally mixed with G.N. 25, from ~~the~~ <sup>10<sup>2</sup></sup> up  
 the Bl. New Town - there OK 367 was nearly the  
 same as G.N. 25 ( $\approx$  S. 497) but with more of Nit. 2. libt. 8cm.

S

637 (New Zealand - c.t.)  
*arlaudianum* (as in 630) *Coscinodiscus* -  
*cocconae* - *caeruleus*.  
*Aimantidium*?  $n \pm 10^{\circ}$   
 nearly, but not quite the same as 630 <sup>more</sup> ~~not~~  
 autocad. here.

638 (New Zealand c.t.)  
*cocconae* n.sp.  $n \pm 10^{\circ}$   
*Coscinodiscus*

639 (Lyamouth - A. Taylor no. 34)  
*Rav. pectinaria* ~~estab.~~ *Bailey* in  
 monograph for S.V. of New Brit  
 (See # 630 for better supply)

S

640 (Stomach of oyster-hull  
Rhizosolenia styliformis, etc.)

641 (New Zealand, mountains 150 m. from Christchurch S.W.)  
Epithemial sp.  
Gonyphorme  
(most like same as 5.595).

642 (Side of Loch Tay - Putherton - G. Norm. no 70)  
Gonyphorme ventricosa, geminata

see 365 254

643 (Westdale Beach, etc., etc.)

64

~~644~~ <sup>651</sup> Bidston marsh - J.C. Aug 58  
Fossil capitate or <sup>645</sup> gracile (middle of rock)  
~~651~~ Southport sand (J.C. Aug 58) m.  
Haworthia pulchella var (now Haworthia mucronata  
var. pulchella) <sup>new synonymy</sup>  
Southport sand R. Sept 58  
10 miles N.E. of 29 miles to the N.E.

Emptied

Matale sand, New Zealand May 1901  
R. L. M. 565  
Pecten affinis.

Spithemis contracta Dub (n.n. Sp.). Mastogloia lancea  
W. Nitze. on 18650. Surinam Brightwell var?

30. 3. 39

Nitzschia! (newly known  
ad linear) var. Brightwell? many in last

651 / ~~Pedicularis~~ See last page  
for who? J. S. 1865. 677

652 Wall of rock tank 8 m. S. 21. 1. 50

Leptothrix minuta (Oppenb.) 397 & 356  
leaves strongly developed from the bottom of the umbelifer

653 D. S. 1865. 678

654 200' above H. Station <sup>Path</sup>  
Baccharis form. ligustrina  
Stream at Church <sup>Street</sup> - ~~Hill~~ - L. M. B.  
bottom. Yellow, yellowish, reddish, orange,

656 Sweet Heart Rock 20.

7058 / G. N. 1865. 678  
Pteridium aquilinum?  
Pteridium aquilinum?  
71 Sampson Bay  
Ferns galore, slugs, by sea fo-

223 (Green Park, January 1855 D. Gray C. 132)  
"Achnanthidium lanceolatum" (n. s.)

224 (Haverfordwest T. Okeden OK. 36)  
Nauclea cryptoceras var angustata?

225 (Haverfordwest T. Okeden OK. 78)  
Meniscium circulare; Achnanthidium lan-  
ceolatum.

erect. First flowering on slides  
and then mounted in balsam

5

- 226 (Dorset thin L. Kedron - B.K. 64)  
*Pleurosigma attenuatum*, lacustr. Naviculae & hypochalata inflata, producta, cuspidata, amphistomata (ren.), Pinnul. viridula, oblonga, Taurinae Phaner., gracilis, N. A. L. linearis & N. medii. Pinn. dim.)

- 227 (Canty Peak Coopers Past. : )  
 > *Orthosia horvathi* (as in Tom Brady deposit)  
*Spirula elegans*, nobilis (Seery) Epithem. Ryndum  
*Staurostoma acutum*

- 228 (Canty Peak, ~~nearer~~<sup>all</sup> ~~down~~ down) )  
 Orthosia (Lever. nobilis  
 (both scars) Epithem. Ryndum

- 5.659 (South Park Sand, light) Corals - Tr. natalensis (Haw) same as from Crofton  
 5.660 (Linen Water, Park, corals) Pinn. late, viridula B  
 5.661 (Swanenden dock - off. 436") amphora Pinn. gracilis  
 Test. apic. in - br. 162P.  
 5.662 (Valley & meadow, Reward off. 630) Amphor. minutissima?  
 and - for Corals der 570  
 5.663 (Swanenden 638") Schizog. - ramose? punctata  
 bivalve der 638" S. Schizog. - ramose?  
 5.664 (Halterprice Hill G.H. 117") Nitz tenuis Cyath. solida  
 Clew. attenuata, Stauro. gracilis - very clear  
 5.665 Hornsea Peak - deposit no Hull - S. H. 164  
 (Moss. punctata, Cyath. solida, Stauro. acuta / non  
 Test. Test. & Grains.  
 5.666 Mull Pond, oyster bed, N. York G.N. - a 81 m.s.t.  
 marine. like sand w/ low Recent species found  
 5.667 Swanenden dock (off. 436") fragment of a large marine  
 bivalve (Spirula?) (Glycimeris) broken and irregular, as  
 Test. Test. & Grains. Stauro. sp. and Cyath. sp.  
 and a small coral (as Hammaria sp. like Hammaria  
 undulata) Part - broken and flat - not whole  
 marine. like sand w/ low Recent species found  
 5.668 Swanenden dock (off. 436") fragment of a large marine  
 bivalve (Spirula?) (Glycimeris) broken and irregular, as  
 Test. Test. & Grains. Stauro. sp. and Cyath. sp.  
 5.669 Swanenden dock (off. 436") fragment of a large marine  
 bivalve (Spirula?) (Glycimeris) broken and irregular, as  
 Test. Test. & Grains. Stauro. sp. and Cyath. sp.  
 5.670 Swanenden dock (off. 436") fragment of a large marine  
 bivalve (Spirula?) (Glycimeris) broken and irregular, as  
 Test. Test. & Grains. Stauro. sp. and Cyath. sp.  
 5.671 ~~Glycimeris~~ Polydella (G.M. Bowring) Campanularia  
 (Cerithidea) (H. H. Stearns) Planorbis, leucostoma, Cucumerulus  
 5.672 (Bromwold Heng, Cambria E.M. Bowring) ~~Glycimeris~~ Semifusca?  
 cyathula reticulata?  
 5.673 Wall at end of first flat Ruthland S.M. Bowring  
 corals! Syphona minima?  
 5.674 (South Hinkley, off. 9. H. Bowring) a gulf near  
 sea sand Cucumerulus & leucostoma ad. Planorbis  
 cerithidea acutula, etc.  
 5.675 (Riverside, Ripon - in meadow  
 Syphona dilatata, Syphona minima, Cerithidea acutula,  
 5.676 very light of 5.668 (w/ greenish "mud" bottom)

- S. 677 - Sennet Land, Jura, Scotland. S. 681  
 S. 678 (Stony cliff, Grindavik) N. 106  
 S. 678 (Cape Horn, Aug 1897) Synchropterus  
 Joseph Beale's Gran. moulins, French Polynesia  
 Pines, 3000 ft. - 10000 ft. - 11000 ft.  
 S. 680 West Hill, Duncansfield, Arthur's Seat / Edinburgh, Scotland  
 King's Mill - Broughton. Middle & large size  
 fish in both - very  
 S. 681 Foreton Rock, Cuckmere T. Cuck.  
 about 100 feet above sea level  
 S. 682 new bright sand, like  
 andักษ์ dappled rock  
 S. 683 (Sennet Land, N. 106) Pleuro. column  
 S. 684 (Hull, 1901) S. N. 106. Same  
 S. 685 (N. 106) S. N. 128. Gramm. column,  
 flatellus  
 S. 686 (Northern) S. 129 - Synchropterus  
 megalopter - like leaf on bottom?  
 S. 687 (Northern, Norway) N. 130. Synchropterus column  
 S. 688 (Ditch at Kinnordy, Orkney Islands, October 9, 1901)  
 in the Glen Firth, Orkney Islands. Epithem gill  
 some marked S. 688<sup>2</sup> were from the headland, others  
 after winds, when more  
 S. 689 Barrow castle, N.H. a few feet from among  
 many Ophiodes - Triceratinae - to an  
 extent than around Skye?  
 S. 690 (Orkney) Acanthoceras. (in) Phryg. Delt.  
 S. 691 (or) Rhynch. Delt. in Red M. Bay  
 S. 692 (or) Chrysanthemum (with crabs) Name - Syn. or?  
 (small, fine thorns)

S. 693 (or) Rhynch. Delt. in Red M. Bay  
 (likely the right side of the bay of the  
 above S. 690 & 691 - 3000 ft. high)  
 small, fine thorns & broken

S. 694

	Med. Delt.
10	+116 ph
12	25
14	6
16	3
18	317
20	320
22	345 ph
24	373
26	383
28	502
30	25
32	405
34	472
36	473
38	471
40	489
42	495
44	513
46	528
48	5
50	+666 quite rare

local S. 682  
 filled with brownish  
 surface No. 2

in fly S. 9 - 32  
 etc.

S. 685

S. 686

S. 686 the following  
 a number of them  
 green

Those marked X  
 may be certainly  
 Northern way.